

Sta 711: Homework 7

Almost-sure Convergence

1. Let $\{X_n\}$ be a monotonically increasing sequence of RVs such that $X_n \rightarrow X$ in probability (*pr.*). Show that $X_n \rightarrow X$ almost surely (*a.s.*)
2. Let $\{X_n\}$ be any sequence of RVs. Show that $X_n \rightarrow X$ *a.s.* if and only if

$$\sup_{k \geq n} |X_k - X| \rightarrow 0 \quad \text{pr.}$$

3. Let $\{X_n\}$ be an arbitrary sequence of RVs and set $S_n := \sum_{i=1}^n X_i$. Show that $X_n \rightarrow 0$ *a.s.* implies that $S_n/n \rightarrow 0$ *a.s.*

In-probability Convergence

4. Let $\{X_n\} \subset L_2$ be independent and identically distributed. For each $\delta > 0$ show that $n\mathbb{P}[|X_1| > \delta\sqrt{n}] \rightarrow 0$. Use this to show that the maximum $\sqrt{\sum_{i=1}^n |X_i|}/\sqrt{n} \rightarrow 0$ *pr.* Thus, the maximum of n iid L_2 random variables grows slower than \sqrt{n} .
5. For random variables X, Y define

$$\rho(X, Y) := \mathbb{E} \left\{ \frac{|X - Y|}{1 + |X - Y|} \right\}$$

The function ρ is a metric (you do *not* have to prove that), *i.e.*, it's non-negative, symmetric, satisfies the triangle inequality, and vanishes if and only if $X = Y$ *a.s.* Show that $X_n \rightarrow X$ *pr.* if and only if $\rho(X_n, X) \rightarrow 0$. Thus, convergence in probability is metrizable.¹

L_p Convergence

6. Find a sequence of RVs $\{X_n\} \subset L_2$ which converge in L_1 but not in L_2 .
7. Let $(\Omega, \mathcal{F}, \mathbb{P}) := ((0, 1], \mathcal{B}, \lambda)$ be the unit interval with Borel sets and Lebesgue measure and define $X_n(\omega) := \omega^n$ for $n \in \mathbb{N}$, $\omega \in \Omega$. For what $p \in [1, \infty]$, does the sequence $\{X_n\}$ converge in L_p ? To what limit? Explain your answer.
8. Verify Hölder's inequality for $p = 1$, $q = \infty$ and all random variables X, Y :

$$\mathbb{E}|XY| \leq \|X\|_1 \|Y\|_\infty$$

where $\|Y\|_\infty := \sup\{c < \infty : \mathbb{P}[|Y| > c] > 0\}$.

9. Verify Minkowski's inequality for $p = \infty$ and all random variables X, Y :

$$\|X + Y\|_\infty \leq \|X\|_\infty + \|Y\|_\infty$$

¹Many other metrics would work too—like $\mathbb{E}(|X - Y| \wedge 1)$ or $\inf\{\epsilon > 0 : \mathbb{P}[|X - Y| > \epsilon] \leq \epsilon\}$.

Uniform Integrability (UI)

10. Fix $p > 0$ and set $X_n := n^p \mathbf{1}_{\{0 < \omega \leq 1/n\}}$ on $(\Omega, \mathcal{F}, \mathbb{P})$ with $\Omega = (0, 1]$, $\mathcal{F} = \mathcal{B}(\Omega)$, and $\mathbb{P} = \lambda$. Show explicitly that $\{X_n\}$ is UI for $p < 1$ and not for $p \geq 1$, by verifying that $\mathbb{E}[X_n \mathbf{1}_{\{X_n > t\}}]$ converges to zero uniformly as $t \rightarrow \infty$ for $p < 1$ and not for $p \geq 1$.
11. Let $\{X_n\}$ be an iid sequence of L_1 random variables and set $S_n := \sum_{i=1}^n X_i$. Show that the sequence of random variables $\{\bar{X}_n\}$ defined by $\bar{X}_n := S_n/n$ is UI.
12. Let $\{X_n\}$ be iid and L_1 . Show²:

$$\mathbb{P}\left(\lim_{n \rightarrow \infty} \frac{X_n}{n} = 0\right) = 1.$$

13. If $\{X_n\}$ and $\{Y_n\}$ are UI, show that so is $\{X_n + Y_n\}$.
14. Let $\phi(x) \geq 0$ be a nonnegative function which grows faster than x at infinity, *i.e.*, $\phi(x)/x \rightarrow \infty$ as $x \rightarrow \infty$. Let \mathcal{C} be a collection of random variables such that, for some fixed $B < \infty$ and all $Z \in \mathcal{C}$,

$$\mathbb{E}(\phi(|Z|)) \leq B.$$

Show that \mathcal{C} is UI. In particular, any collection of random variables that is bounded uniformly in L_p for some $p > 1$ is also UI.

²Although $\{X_i\}$ are UI, that won't be a factor in solving this problem. Is independence needed?