

Name	Notation	pdf/pmf	Range	Mean μ	Variance σ^2
Beta	$\text{Be}(\alpha, \beta)$	$f(x) = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)}x^{\alpha-1}(1-x)^{\beta-1}$	$x \in (0, 1)$	$\frac{\alpha}{\alpha+\beta}$	$\frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$
Binomial	$\text{Bi}(n, p)$	$f(x) = \binom{n}{x} p^x q^{(n-x)}$	$x \in 0, \dots, n$	$n p$	$n p q \quad (q = 1 - p)$
Exponential	$\text{Ex}(\lambda)$	$f(x) = \lambda e^{-\lambda x}$	$x \in \mathbb{R}_+$	$1/\lambda$	$1/\lambda^2$
Gamma	$\text{Ga}(\alpha, \lambda)$	$f(x) = \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x}$	$x \in \mathbb{R}_+$	α/λ	α/λ^2
Geometric	$\text{Ge}(p)$	$f(x) = p q^x$	$x \in \mathbb{Z}_+$	q/p	$q/p^2 \quad (q = 1 - p)$
HyperGeometric	$\text{HG}(n, A, B)$	$f(x) = \binom{A}{x} \binom{B}{n-x} / \binom{A+B}{n}$	$x \in 0, \dots, n$	$n P$	$n P Q \frac{N-n}{N-1} \quad (P = \frac{A}{A+B}, Q = \frac{B}{A+B})$
Logistic	$\text{Lo}(\mu, \beta)$	$f(x) = \frac{e^{-(x-\mu)/\beta}}{\beta[1+e^{-(x-\mu)/\beta}]^2}$	$x \in \mathbb{R}$	μ	$\pi^2 \beta^2 / 3$
Log Normal	$\text{LN}(\mu, \sigma^2)$	$f(x) = \frac{1}{x\sqrt{2\pi\sigma^2}} e^{-(\log x - \mu)^2 / 2\sigma^2}$	$x \in \mathbb{R}_+$	$e^{\mu + \sigma^2/2}$	$e^{2\mu + \sigma^2} (1 - e^{\sigma^2})$
Negative Binomial	$\text{NB}(\alpha, p)$	$f(x) = \binom{-\alpha}{x} p^\alpha (-q)^x$	$x \in \mathbb{Z}_+$	$\alpha q / p$	$\alpha q / p^2 \quad (q = 1 - p)$
Normal	$\text{No}(\mu, \sigma^2)$	$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\mu)^2 / 2\sigma^2}$	$x \in \mathbb{R}$	μ	σ^2
Pareto	$\text{Pa}(\alpha, \beta)$	$f(x) = \beta \alpha^\beta / x^{\beta+1}$	$x \in (\alpha, \infty)$	$\frac{\alpha\beta}{\beta-1}$	$\frac{\alpha^2\beta}{(\beta-1)^2(\beta-2)}$
Poisson	$\text{Po}(\lambda)$	$f(x) = \frac{\lambda^x}{x!} e^{-\lambda}$	$x \in \mathbb{Z}_+$	λ	λ
Snedecor F	$F(\nu_1, \nu_2)$	$f(x) = \frac{\Gamma(\frac{\nu_1+\nu_2}{2})(\nu_1/\nu_2)^{\nu_1/2}}{\Gamma(\nu_1/2) \Gamma(\nu_2/2)} \times x^{\frac{\nu_1-2}{2}} \left[1 + \frac{\nu_1}{\nu_2} x\right]^{-\frac{\nu_1+\nu_2}{2}}$	$x \in \mathbb{R}_+$	$\frac{\nu_2}{\nu_2-2}$	$\left(\frac{\nu_2}{\nu_2-2}\right)^2 \frac{2(\nu_1+\nu_2-2)}{\nu_1(\nu_2-4)}$
Student t	$t(\nu)$	$f(x) = \frac{\Gamma(\frac{\nu+1}{2})}{\Gamma(\frac{\nu}{2})\sqrt{\pi\nu}} [1 + x^2/\nu]^{-(\nu+1)/2}$	$x \in \mathbb{R}$	0	$\nu / (\nu - 2)$
Uniform	$\text{Un}(\alpha, \beta)$	$f(x) = \frac{1}{\beta-\alpha}$	$x \in (\alpha, \beta)$	$\frac{\alpha+\beta}{2}$	$\frac{(\beta-\alpha)^2}{12}$