

# Lecture 15 - Tests of Two Proportions

---

Sta102 / BME102

March 23, 2016

Colin Rundel

## Difference of two proportions

---

## Example - Melting ice cap survey

Scientists predict that global warming may have big effects on the polar regions within the next 100 years. One of the possible effects is that the northern ice cap may completely melt. Would this bother you a great deal, some, a little, or not at all if it actually happened?

- (a) A great deal
- (b) Some
- (c) A little
- (d) Not at all

## Results from the GSS & Duke

The GSS asks this question, below is the distribution of responses from the 2010 survey:

A great deal	454
Not a great deal	226
Total	680

## Results from the GSS & Duke

The GSS asks this question, below is the distribution of responses from the 2010 survey:

A great deal	454
Not a great deal	226
Total	680

The same question was asked of 88 Duke students, of which 56 said it would bother them a great deal.

We will collapse the data such that we consider only the responses of a great deal and its compliment, not a great deal.

## Collapsed Results

	US	Duke	Total
A great deal	454	56	510
Not a great deal	226	32	258
Total	680	88	768

This is an example of a contingency table (specifically a 2 x 2 contingency table).

## Collapsed Results

	US	Duke	Total
A great deal	454	56	510
Not a great deal	226	32	258
Total	680	88	768

This is an example of a contingency table (specifically a 2 x 2 contingency table).

We are interested in comparing proportion of Duke students who say it would both them a great deal ( $p_{GD|Duke} = 56/88$ ) to the proportion of all Americans who say it would both them a great deal ( $p_{GD|US} = 454/680$ ).

## Condition on what?

Knowing which of the two variables to condition on can be tricky some times.

Ask yourself - which of the two variables is most likely the dependent variable ( $y$ ) and which is most likely the independent variable ( $x$ ). In other words, changes in  $x$  should *cause* changes in  $y$  (not the other way around).

Once we know this then the two proportions of interest are:

$$p_{y_1|x_1} \quad \text{and} \quad p_{y_1|x_2}$$



## Parameter and point estimate

- *Parameter of interest*: Difference between the proportions of *all* Duke students and *all* Americans who would be bothered a great deal by the northern ice cap melting.

$$P_{GD|Duke} - P_{GD|US}$$

## Parameter and point estimate

- *Parameter of interest*: Difference between the proportions of *all* Duke students and *all* Americans who would be bothered a great deal by the northern ice cap melting.

$$P_{GD|Duke} - P_{GD|US}$$

- *Point estimate*: Difference between the proportions of *sampled* Duke students and *sampled* Americans who would be bothered a great deal by the northern ice cap melting.

$$\hat{P}_{GD|Duke} - \hat{P}_{GD|US}$$

## Inference for comparing proportions

The details for inference are the same as what we've seen previously,

## Inference for comparing proportions

The details for inference are the same as what we've seen previously,

- CI: *point estimate  $\pm$  critical value  $\times$  std error*

# Inference for comparing proportions

The details for inference are the same as what we've seen previously,

- CI:  $\text{point estimate} \pm \text{critical value} \times \text{std error}$
- HT:  $\text{Test Statistic} = \frac{\text{point estimate} - \text{null value}}{\text{std error}}$ , find appropriate p-value using sampling distribution.

# Inference for comparing proportions

The details for inference are the same as what we've seen previously,

- CI:  $\text{point estimate} \pm \text{critical value} \times \text{std error}$
- HT:  $\text{Test Statistic} = \frac{\text{point estimate} - \text{null value}}{\text{std error}}$ , find appropriate p-value using sampling distribution.
- We just need to figure out the appropriate sampling distribution and its parameters..

## Sampling Distribution

Last time we saw that the sampling distribution for  $\hat{p}$  is a normal with mean  $p$  and variance  $p(1 - p)/n$ .

## Sampling Distribution

Last time we saw that the sampling distribution for  $\hat{p}$  is a normal with mean  $p$  and variance  $p(1 - p)/n$ .

We can combine that result with the approach we used for the test of two means to find the distribution of  $\hat{p}_1 - \hat{p}_2$

$$\hat{p}_1 - \hat{p}_2 \sim N(\mu = E(\hat{p}_1 - \hat{p}_2), \sigma^2 = \text{Var}(\hat{p}_1 - \hat{p}_2))$$



## Sampling Distribution

Last time we saw that the sampling distribution for  $\hat{p}$  is a normal with mean  $p$  and variance  $p(1-p)/n$ .

We can combine that result with the approach we used for the test of two means to find the distribution of  $\hat{p}_1 - \hat{p}_2$

$$\hat{p}_1 - \hat{p}_2 \sim N(\mu = E(\hat{p}_1 - \hat{p}_2), \sigma^2 = \text{Var}(\hat{p}_1 - \hat{p}_2))$$

$$\begin{aligned} E(\hat{p}_1 - \hat{p}_2) &= E(\hat{p}_1) - E(\hat{p}_2) & \text{Var}(\hat{p}_1 - \hat{p}_2) &= \text{Var}(\hat{p}_1) + \text{Var}(\hat{p}_2) \\ &= p_1 - p_2 & &= \frac{p_1(1-p_1)}{n_1} + \frac{p_2(1-p_2)}{n} \end{aligned}$$

## Sampling Distribution

Last time we saw that the sampling distribution for  $\hat{p}$  is a normal with mean  $p$  and variance  $p(1-p)/n$ .

We can combine that result with the approach we used for the test of two means to find the distribution of  $\hat{p}_1 - \hat{p}_2$

$$\hat{p}_1 - \hat{p}_2 \sim N(\mu = E(\hat{p}_1 - \hat{p}_2), \sigma^2 = \text{Var}(\hat{p}_1 - \hat{p}_2))$$

$$\begin{aligned} E(\hat{p}_1 - \hat{p}_2) &= E(\hat{p}_1) - E(\hat{p}_2) & \text{Var}(\hat{p}_1 - \hat{p}_2) &= \text{Var}(\hat{p}_1) + \text{Var}(\hat{p}_2) \\ &= p_1 - p_2 & &= \frac{p_1(1-p_1)}{n_1} + \frac{p_2(1-p_2)}{n} \end{aligned}$$

Note - as with the test of two means, this result requires that  $\hat{p}_1$  and  $\hat{p}_2$  are independent.

## Conditions for CI for the difference of two proportions

1. *Independence within groups:*

- The US group is sampled randomly and we're assuming that the Duke group represents a random sample as well.

# Conditions for CI for the difference of two proportions

## 1. *Independence within groups:*

- The US group is sampled randomly and we're assuming that the Duke group represents a random sample as well.
- $n_{Duke} < 10\%$  of all Duke students and  $680 < 10\%$  of all Americans.

# Conditions for CI for the difference of two proportions

## 1. *Independence within groups:*

- The US group is sampled randomly and we're assuming that the Duke group represents a random sample as well.
- $n_{Duke} < 10\%$  of all Duke students and  $680 < 10\%$  of all Americans.

We can assume that the attitudes of Duke students in the sample are independent of each other, and attitudes of US residents in the sample are independent of each other as well.

# Conditions for CI for the difference of two proportions

## 1. *Independence within groups:*

- The US group is sampled randomly and we're assuming that the Duke group represents a random sample as well.
- $n_{Duke} < 10\%$  of all Duke students and  $680 < 10\%$  of all Americans.

We can assume that the attitudes of Duke students in the sample are independent of each other, and attitudes of US residents in the sample are independent of each other as well.

- ## 2. *Independence between groups:* The sampled Duke students and the US residents are independent of each other.

# Conditions for CI for the difference of two proportions

## 1. *Independence within groups:*

- The US group is sampled randomly and we're assuming that the Duke group represents a random sample as well.
- $n_{Duke} < 10\%$  of all Duke students and  $680 < 10\%$  of all Americans.

We can assume that the attitudes of Duke students in the sample are independent of each other, and attitudes of US residents in the sample are independent of each other as well.

- ## 2. *Independence between groups:* The sampled Duke students and the US residents are independent of each other.
- ## 3. *Success-failure:*

At least 10 observed successes and 10 observed failures in *both* groups.

## CI for difference of proportions

Construct a 95% confidence interval for the difference between the proportions of Duke students and Americans who would be bothered a great deal by the melting of the northern ice cap ( $p_{GD|Duke} - p_{GD|US}$ ).

	Duke	US
A great deal	56	454
Not a great deal	32	226
Total	88	680



## CI for difference of proportions

Construct a 95% confidence interval for the difference between the proportions of Duke students and Americans who would be bothered a great deal by the melting of the northern ice cap ( $p_{GD|Duke} - p_{GD|US}$ ).

	Duke	US
A great deal	56	454
Not a great deal	32	226
Total	88	680

$$\hat{p}_{GD|Duke} = 56/88 = 0.636$$

$$\hat{p}_{GD|US} = 454/680 = 0.668$$

## CI for difference of proportions

Construct a 95% confidence interval for the difference between the proportions of Duke students and Americans who would be bothered a great deal by the melting of the northern ice cap ( $p_{GD|Duke} - p_{GD|US}$ ).

	Duke	US
A great deal	56	454
Not a great deal	32	226
Total	88	680

$$\hat{p}_{GD|Duke} = 56/88 = 0.636$$

$$\hat{p}_{GD|US} = 454/680 = 0.668$$

$$\begin{aligned} SE &\approx \sqrt{\frac{\hat{p}_{GD|Duke}(1 - \hat{p}_{GD|Duke})}{n_{Duke}} + \frac{\hat{p}_{GD|US}(1 - \hat{p}_{GD|US})}{n_{US}}} \\ &= \sqrt{\frac{0.636(1 - 0.636)}{88} + \frac{0.668(1 - 0.668)}{680}} = 0.0537 \end{aligned}$$

## CI for difference of proportions, cont.

$$\hat{p}_{GD|Duke} = 0.636$$

$$\hat{p}_{GD|US} = 0.668$$

$$SE = 0.0537$$

## CI for difference of proportions, cont.

$$\hat{p}_{GD|Duke} = 0.636$$

$$\hat{p}_{GD|US} = 0.668$$

$$SE = 0.0537$$

$$CI = PE \pm CV \times SE$$

$$= (\hat{p}_{GD|Duke} - \hat{p}_{GD|US}) \pm Z^* \times \sqrt{\frac{\hat{p}_{GD|Duke}(1 - \hat{p}_{GD|Duke})}{n_{Duke}}}$$

$$= (0.636 - 0.668) \pm 1.96 \times 0.0537$$

$$= (-0.138, 0.074)$$

## CI for difference of proportions, cont.

$$\hat{p}_{GD|Duke} = 0.636$$

$$\hat{p}_{GD|US} = 0.668$$

$$SE = 0.0537$$

$$CI = PE \pm CV \times SE$$

$$= (\hat{p}_{GD|Duke} - \hat{p}_{GD|US}) \pm Z^* \times \sqrt{\frac{\hat{p}_{GD|Duke}(1 - \hat{p}_{GD|Duke})}{n_{Duke}}}$$

$$= (0.636 - 0.668) \pm 1.96 \times 0.0537$$

$$= (-0.138, 0.074)$$

What conclusion should we draw here?

## Hypotheses for testing the difference of two proportions

Just like the other hypothesis tests we have seen thus far, we formulate our null and alternative hypotheses for testing if the proportion of all Duke students who would be bothered a great deal by the melting of the northern ice cap differs from the proportion of all Americans who do as follows,

$$H_0 : p_{GD|Duke} = p_{GD|US} \quad \Rightarrow \quad p_{GD|Duke} - p_{GD|US} = 0$$

$$H_A : p_{GD|Duke} \neq p_{GD|US} \quad \Rightarrow \quad p_{GD|Duke} - p_{GD|US} \neq 0$$

## Flashback to working with one proportion

When constructing a confidence interval for a population proportion, we check if the *observed* number of successes and failures are at least 10.

$$n\hat{p} \geq 10$$

$$n(1 - \hat{p}) \geq 10$$

## Flashback to working with one proportion

When constructing a confidence interval for a population proportion, we check if the *observed* number of successes and failures are at least 10.

$$n\hat{p} \geq 10 \qquad n(1 - \hat{p}) \geq 10$$

When conducting a hypothesis test for a population proportion, we check if the *expected* number of successes and failures are at least 10.

$$np_0 \geq 10 \qquad n(1 - p_0) \geq 10$$



## A slight wrinkle ...

In setting the null hypothesis for comparing two proportions we haven't fixed either  $p_{GD|Duke}$  or  $p_{GD|US}$  - instead we have fixed their difference.

## A slight wrinkle ...

In setting the null hypothesis for comparing two proportions we haven't fixed either  $p_{GD|Duke}$  or  $p_{GD|US}$  - instead we have fixed their difference.

As such, we don't have a specific null value we can use to calculate the *expected* number of successes and failures in each group or the standard error. So, we know the following

## A slight wrinkle ...

In setting the null hypothesis for comparing two proportions we haven't fixed either  $p_{GD|Duke}$  or  $p_{GD|US}$  - instead we have fixed their difference.

As such, we don't have a specific null value we can use to calculate the *expected* number of successes and failures in each group or the standard error. So, we know the following

$$p_{GD|Duke} = p_{GD|US}$$

$$p_{GD|Duke} = ?$$

$$p_{GD|US} = ?$$

## A slight wrinkle ...

In setting the null hypothesis for comparing two proportions we haven't fixed either  $p_{GD|Duke}$  or  $p_{GD|US}$  - instead we have fixed their difference.

As such, we don't have a specific null value we can use to calculate the *expected* number of successes and failures in each group or the standard error. So, we know the following

$$p_{GD|Duke} = p_{GD|US}$$

$$p_{GD|Duke} = ?$$

$$p_{GD|US} = ?$$

Does this null give us any additional useful information?

# Proportions and Probabilities

Think about the sample proportions as probabilities, what does it mean if

$$P(GD|Duke) = P(GD|US)$$

# Proportions and Probabilities

Think about the sample proportions as probabilities, what does it mean if

$$P(GD|Duke) = P(GD|US)$$

If these two probabilities are equal then global warming concern is *independent* of the Duke vs. US grouping. Which means that,

$$P(GD|Duke) = P(GD|US) = P(GD)$$

# Pooling

As such, our null hypothesis is equivalent to claiming that our two categorical variables are independent. So when conducting the hypothesis test we assume the null hypothesis to be true, which means we must also assume that the two variables are independent.

Under the assumption of independence our best guess for both  $p_{GD|Duke}$  and  $p_{GD|US}$  will be  $\hat{p}_{GD}$ , which is the sample proportion of *all* respondents (from Duke or US) who answered “A great deal”.

We call this value  $\hat{p}_{pooled}$ ,

$$\hat{p}_{pooled} = \frac{\# \text{ of successes in 1} + \# \text{ of successes in 2}}{n_1 + n_2} = \frac{n_1 \hat{p}_1 + n_2 \hat{p}_2}{n_1 + n_2}$$

## Pooled estimate of a proportion

Calculate the estimated pooled proportion of Duke students and Americans who would be bothered a great deal by the melting of the northern ice cap.

	Duke	US	Total
A great deal	56	454	510
Not a great deal	32	226	258
Total	88	680	788

$$\hat{p}_{pooled} = \frac{56 + 454}{88 + 680} = \frac{510}{788} = 0.664$$



## Pooled estimate of a proportion

Calculate the estimated pooled proportion of Duke students and Americans who would be bothered a great deal by the melting of the northern ice cap.

	Duke	US	Total
A great deal	56	454	510
Not a great deal	32	226	258
Total	88	680	788

$$\hat{p}_{pooled} = \frac{56 + 454}{88 + 680} = \frac{510}{788} = 0.664$$

Which sample proportion ( $\hat{p}_{GD|Duke}$  or  $\hat{p}_{GD|US}$ ) is closer to the pooled estimate? Why?

## Implications for the SE

Under the null hypothesis we are stating that  $p_1 = p_2$  which does not uniquely identify either  $p_1$  or  $p_2$ . Therefore we are using the pooled proportion ( $\hat{p}$ ) as our best guess for  $p_1$  and  $p_2$  under the null hypothesis.

For a *confidence interval* we use  $\hat{p}_1$  and  $\hat{p}_2$  to approximate for  $p_1$  and  $p_2$

$$SE = \sqrt{\frac{p_1(1-p_1)}{n_1} + \frac{p_2(1-p_2)}{n_2}} \approx \sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n_1} + \frac{\hat{p}_2(1-\hat{p}_2)}{n_2}}$$

While for a *hypothesis test* we use  $\hat{p}_{pooled}$  to approximate for  $p_1$  and  $p_2$

$$\begin{aligned} SE &= \sqrt{\frac{p_1(1-p_1)}{n_1} + \frac{p_2(1-p_2)}{n_2}} \approx \sqrt{\frac{\hat{p}_p(1-\hat{p}_p)}{n_1} + \frac{\hat{p}_p(1-\hat{p}_p)}{n_2}} \\ &= \sqrt{\hat{p}_p(1-\hat{p}_p) \left( \frac{1}{n_1} + \frac{1}{n_2} \right)} \end{aligned}$$

## HT for comparing proportions

Do these data suggest that the proportion of all Duke students who would be bothered a great deal by the melting of the northern ice cap differs from the proportion of all Americans who do?

$$\hat{p}_{pooled} = 0.664, \quad n_1 = 88, \quad n_2 = 680$$

## HT for comparing proportions

Do these data suggest that the proportion of all Duke students who would be bothered a great deal by the melting of the northern ice cap differs from the proportion of all Americans who do?

$$\hat{p}_{pooled} = 0.664, \quad n_1 = 88, \quad n_2 = 680$$

$$SE = \sqrt{\hat{p}_p(1 - \hat{p}_p) \left( \frac{1}{n_1} + \frac{1}{n_2} \right)} = \sqrt{0.664(1 - 0.664) \left( \frac{1}{88} + \frac{1}{680} \right)} = 0.0535$$

## HT for comparing proportions

Do these data suggest that the proportion of all Duke students who would be bothered a great deal by the melting of the northern ice cap differs from the proportion of all Americans who do?

$$\hat{p}_{pooled} = 0.664, \quad n_1 = 88, \quad n_2 = 680$$

$$SE = \sqrt{\hat{p}_p(1 - \hat{p}_p) \left( \frac{1}{n_1} + \frac{1}{n_2} \right)} = \sqrt{0.664(1 - 0.664) \left( \frac{1}{88} + \frac{1}{680} \right)} = 0.0535$$

$$Z = \frac{(56/88 - 454/680) - 0}{0.0535} = -0.59$$

## HT for comparing proportions

Do these data suggest that the proportion of all Duke students who would be bothered a great deal by the melting of the northern ice cap differs from the proportion of all Americans who do?

$$\hat{p}_{pooled} = 0.664, \quad n_1 = 88, \quad n_2 = 680$$

$$SE = \sqrt{\hat{p}_p(1 - \hat{p}_p) \left( \frac{1}{n_1} + \frac{1}{n_2} \right)} = \sqrt{0.664(1 - 0.664) \left( \frac{1}{88} + \frac{1}{680} \right)} = 0.0535$$

$$Z = \frac{(56/88 - 454/680) - 0}{0.0535} = -0.59$$

$$p\text{-value} = P(Z < -0.59 \text{ or } Z > 0.59)$$

$$= 0.277 + 0.277 = 0.555$$

## HT and CI agreement

Confidence interval:

$$CI = (-0.138, 0.074)$$

Hypothesis test:

$$H_0 : p_{GD|Duke} = p_{GD|US}$$

$$Z = -0.59$$

$$H_A : p_{GD|Duke} \neq p_{GD|US}$$

$$p\text{-value} = 0.555$$

Do the results of the Confidence interval and hypothesis test agree? Do they necessarily have to agree?

## Picking successes?

What would happen to our analysis if we had picked “Not a great deal”?

	Duke	US	Total
A great deal	56	454	510
Not a great deal	32	226	258
Total	88	680	788



## Picking successes?

What would happen to our analysis if we had picked “Not a great deal”?

	Duke	US	Total
A great deal	56	454	510
Not a great deal	32	226	258
Total	88	680	788

$$H_0 : p_{NGD|Duke} = p_{NGD|US}$$

$$H_0 : p_{NGD|Duke} \neq p_{NGD|US}$$

$$\hat{p}_{pooled} = \frac{32 + 226}{88 + 680} = \frac{258}{788} = 0.336$$

## Picking successes?

What would happen to our analysis if we had picked “Not a great deal”?

	Duke	US	Total
A great deal	56	454	510
Not a great deal	32	226	258
Total	88	680	788

$$H_0 : p_{NGD|Duke} = p_{NGD|US}$$

$$H_0 : p_{NGD|Duke} \neq p_{NGD|US}$$

$$\hat{p}_{pooled} = \frac{32 + 226}{88 + 680} = \frac{258}{788} = 0.336$$

$$SE = \sqrt{0.336(1 - 0.336) \left( \frac{1}{88} + \frac{1}{680} \right)} = 0.0535$$

## Picking successes?

What would happen to our analysis if we had picked “Not a great deal”?

	Duke	US	Total
A great deal	56	454	510
Not a great deal	32	226	258
Total	88	680	788

$$H_0 : p_{NGD|Duke} = p_{NGD|US}$$

$$H_0 : p_{NGD|Duke} \neq p_{NGD|US}$$

$$\hat{p}_{pooled} = \frac{32 + 226}{88 + 680} = \frac{258}{788} = 0.336$$

$$SE = \sqrt{0.336(1 - 0.336) \left( \frac{1}{88} + \frac{1}{680} \right)} = 0.0535$$

$$Z = \frac{(32/88 - 226/680) - 0}{0.0535} = 0.585$$

$$\begin{aligned} \text{p-value} &= P(Z < -0.59 \text{ or } Z > 0.59) \\ &= 0.277 + 0.277 = 0.555 \end{aligned}$$

## Swapping dependent and independent variables?

What would happen to our analysis if we had swapped our independent and dependent variable?

	Duke	US	Total
A great deal	56	454	510
Not a great deal	32	226	258
Total	88	680	788

## Swapping dependent and independent variables?

What would happen to our analysis if we had swapped our independent and dependent variable?

	Duke	US	Total
A great deal	56	454	510
Not a great deal	32	226	258
Total	88	680	788

$$H_0 : p_{Duke|GD} = p_{Duke|NGD}$$

$$H_0 : p_{Duke|GD} \neq p_{Duke|NGD}$$

$$\hat{p}_{pooled} = \frac{56 + 32}{510 + 258} = \frac{88}{788} = 0.115$$

## Swapping dependent and independent variables?

What would happen to our analysis if we had swapped our independent and dependent variable?

	Duke	US	Total
A great deal	56	454	510
Not a great deal	32	226	258
Total	88	680	788

$$H_0 : p_{Duke|GD} = p_{Duke|NGD}$$

$$H_0 : p_{Duke|GD} \neq p_{Duke|NGD}$$

$$\hat{p}_{pooled} = \frac{56 + 32}{510 + 258} = \frac{88}{788} = 0.115$$

$$SE = \sqrt{0.115(1 - 0.115) \left( \frac{1}{510} + \frac{1}{258} \right)} = 0.0241$$

## Swapping dependent and independent variables?

What would happen to our analysis if we had swapped our independent and dependent variable?

	Duke	US	Total
A great deal	56	454	510
Not a great deal	32	226	258
Total	88	680	788

$$H_0 : p_{Duke|GD} = p_{Duke|NGD}$$

$$H_0 : p_{Duke|GD} \neq p_{Duke|NGD}$$

$$\hat{p}_{pooled} = \frac{56 + 32}{510 + 258} = \frac{88}{788} = 0.115$$

$$SE = \sqrt{0.115(1 - 0.115) \left( \frac{1}{510} + \frac{1}{258} \right)} = 0.0241$$

$$Z = \frac{(56/510 - 32/258) - 0}{0.0241} = 0.59$$

$$\begin{aligned} \text{p-value} &= P(Z < -0.59 \text{ or } Z > 0.59) \\ &= 0.2775 + 0.2775 = 0.555 \end{aligned}$$

What is the power of our hypothesis test to detect a difference of 0.1?

*Step 0:*  $H_0 : p_1 = p_2$     $H_A : p_1 \neq p_2$     $\alpha = 0.05$     $\delta = 0.1$     $SE = 0.0535$    power = ?



What is the power of our hypothesis test to detect a difference of 0.1?

*Step 0:*  $H_0 : p_1 = p_2$   $H_A : p_1 \neq p_2$   $\alpha = 0.05$   $\delta = 0.1$   $SE = 0.0535$  power = ?

*Step 1:* Find  $\hat{p}_1 - \hat{p}_2$  such that we reject  $H_0$ .

What is the power of our hypothesis test to detect a difference of 0.1?

*Step 0:*  $H_0 : p_1 = p_2$   $H_A : p_1 \neq p_2$   $\alpha = 0.05$   $\delta = 0.1$   $SE = 0.0535$  power = ?

*Step 1:* Find  $\hat{p}_1 - \hat{p}_2$  such that we reject  $H_0$ .

$$P(Z < -z \text{ or } Z > z) < 0.05 \Rightarrow z > 1.96$$

What is the power of our hypothesis test to detect a difference of 0.1?

*Step 0:*  $H_0 : p_1 = p_2$   $H_A : p_1 \neq p_2$   $\alpha = 0.05$   $\delta = 0.1$   $SE = 0.0535$  power = ?

*Step 1:* Find  $\hat{p}_1 - \hat{p}_2$  such that we reject  $H_0$ .

$$P(Z < -z \text{ or } Z > z) < 0.05 \Rightarrow z > 1.96$$

$$\frac{(\hat{p}_1 - \hat{p}_2) - 0}{0.0535} < -1.96 \quad \text{or} \quad \frac{(\hat{p}_1 - \hat{p}_2) - 0}{0.0535} > 1.96$$

What is the power of our hypothesis test to detect a difference of 0.1?

*Step 0:*  $H_0 : p_1 = p_2$   $H_A : p_1 \neq p_2$   $\alpha = 0.05$   $\delta = 0.1$   $SE = 0.0535$  power = ?

*Step 1:* Find  $\hat{p}_1 - \hat{p}_2$  such that we reject  $H_0$ .

$$P(Z < -z \text{ or } Z > z) < 0.05 \Rightarrow z > 1.96$$

$$\frac{(\hat{p}_1 - \hat{p}_2) - 0}{0.0535} < -1.96 \quad \text{or} \quad \frac{(\hat{p}_1 - \hat{p}_2) - 0}{0.0535} > 1.96$$

$$\hat{p}_1 - \hat{p}_2 < -0.105 \quad \text{or} \quad \hat{p}_1 - \hat{p}_2 > 0.105$$

What is the power of our hypothesis test to detect a difference of 0.1?

*Step 0:*  $H_0 : p_1 = p_2$   $H_A : p_1 \neq p_2$   $\alpha = 0.05$   $\delta = 0.1$   $SE = 0.0535$  power = ?

*Step 1:* Find  $\hat{p}_1 - \hat{p}_2$  such that we reject  $H_0$ .

$$P(Z < -z \text{ or } Z > z) < 0.05 \Rightarrow z > 1.96$$

$$\frac{(\hat{p}_1 - \hat{p}_2) - 0}{0.0535} < -1.96 \quad \text{or} \quad \frac{(\hat{p}_1 - \hat{p}_2) - 0}{0.0535} > 1.96$$

$$\hat{p}_1 - \hat{p}_2 < -0.105 \quad \text{or} \quad \hat{p}_1 - \hat{p}_2 > 0.105$$

*Step 2:* Assume  $p_1 - p_2 = 0 + \delta = 0.1$  - we no longer assume independence, must use  $SE = 0.0537$  from the CI instead.

$$\begin{aligned} & P(\hat{p}_1 - \hat{p}_2 < -0.105 \quad \text{or} \quad \hat{p}_1 - \hat{p}_2 > 0.105) \\ &= P\left(Z < \frac{-0.105 - 0.1}{0.0527}\right) + P\left(Z > \frac{0.105 - 0.1}{0.0527}\right) \\ &= P(Z < -3.88) + P(Z > 0.094) \end{aligned}$$

## Example - Planned Parenthood

---

## Planned Parenthood

A Pew Research poll conducted between September 22-27, 2015 asked 805 randomly sampled Americans (who self identify as a Democrat or Republican) and ask about their party affiliation and whether they think any budget agreement must eliminate or maintain funding for Planned Parenthood. The distribution of their responses is shown below.

	Eliminate	Maintain	Total
Democrat	45	378	423
Republican	277	105	382
Total	322	483	805

Pew Research Center. *Majority Says Any Budget Deal Must Include Planned Parenthood Funding*. Sep 28, 2015. <http://www.people-press.org/2015/09/28/majority->

Is there evidence that a greater percentage of Democrats support maintaining funding for Planned Parenthood than Republicans?



Is there evidence that a greater percentage of Democrats support maintaining funding for Planned Parenthood than Republicans?

$$H_0 : p_{m|D} = p_{m|R}$$

$$H_A : p_{m|D} > p_{m|R}$$

Is there evidence that a greater percentage of Democrats support maintaining funding for Planned Parenthood than Republicans?

$$H_0 : p_{m|D} = p_{m|R}$$

$$H_A : p_{m|D} > p_{m|R}$$

$$\hat{p}_{pooled} = \frac{378 + 105}{423 + 382} = 0.6$$

$$SE = \sqrt{0.6(1 - 0.6) \left( \frac{1}{423} + \frac{1}{382} \right)} = 0.0346$$

Is there evidence that a greater percentage of Democrats support maintaining funding for Planned Parenthood than Republicans?

$$H_0 : p_{m|D} = p_{m|R}$$

$$H_A : p_{m|D} > p_{m|R}$$

$$\hat{p}_{pooled} = \frac{378 + 105}{423 + 382} = 0.6$$

$$SE = \sqrt{0.6(1 - 0.6) \left( \frac{1}{423} + \frac{1}{382} \right)} = 0.0346$$

$$Z = \frac{(378/423 - 105/382) - 0}{0.0346} = 17.88$$

$$p\text{-value} = P(Z > 17.88) \approx 0$$

Is there evidence that a greater percentage of Democrats support maintaining funding for Planned Parenthood than Republicans?

Is there evidence that a greater percentage of Democrats support maintaining funding for Planned Parenthood than Republicans?

$$\hat{p}_{m|D} = 378/423 = 0.894 \quad \hat{p}_{m|R} = 105/382 = 0.275$$

Is there evidence that a greater percentage of Democrats support maintaining funding for Planned Parenthood than Republicans?

$$\hat{p}_{m|D} = 378/423 = 0.894 \quad \hat{p}_{m|R} = 105/382 = 0.275$$

$$SE = \sqrt{\frac{378/423(1 - 378/423)}{423} + \frac{105/382(1 - 105/382)}{382}} = 0.0273$$

## Analysis - CI

Is there evidence that a greater percentage of Democrats support maintaining funding for Planned Parenthood than Republicans?

$$\hat{p}_{m|D} = 378/423 = 0.894 \quad \hat{p}_{m|R} = 105/382 = 0.275$$

$$SE = \sqrt{\frac{378/423(1 - 378/423)}{423} + \frac{105/382(1 - 105/382)}{382}} = 0.0273$$

$$\begin{aligned} CI &= (\hat{p}_{m|D} - \hat{p}_{m|R}) \pm Z^* SE \\ &= (378/423 - 105/382) \pm 1.96 \times 0.0273 \\ &= (0.565, 0.672) \end{aligned}$$

What is the power of our hypothesis test to detect a difference of 0.62?

*Step 0:*  $H_0 : p_1 = p_2$     $H_A : p_1 > p_2$     $\alpha = 0.05$     $\delta = 0.62$     $SE = 0.0346$    power = ?



# Power

What is the power of our hypothesis test to detect a difference of 0.62?

*Step 0:*  $H_0 : p_1 = p_2$   $H_A : p_1 > p_2$   $\alpha = 0.05$   $\delta = 0.62$   $SE = 0.0346$  power = ?

*Step 1:* Find  $\hat{p}_1 - \hat{p}_2$  such that we reject  $H_0$ .

What is the power of our hypothesis test to detect a difference of 0.62?

*Step 0:*  $H_0 : p_1 = p_2$   $H_A : p_1 > p_2$   $\alpha = 0.05$   $\delta = 0.62$   $SE = 0.0346$  power = ?

*Step 1:* Find  $\hat{p}_1 - \hat{p}_2$  such that we reject  $H_0$ .

$$P(Z > z) < 0.05 \Rightarrow z > 1.644$$

What is the power of our hypothesis test to detect a difference of 0.62?

*Step 0:*  $H_0 : p_1 = p_2$   $H_A : p_1 > p_2$   $\alpha = 0.05$   $\delta = 0.62$   $SE = 0.0346$  power = ?

*Step 1:* Find  $\hat{p}_1 - \hat{p}_2$  such that we reject  $H_0$ .

$$P(Z > z) < 0.05 \Rightarrow z > 1.644$$

$$\frac{(\hat{p}_1 - \hat{p}_2) - 0}{0.0346} > 1.644$$

What is the power of our hypothesis test to detect a difference of 0.62?

*Step 0:*  $H_0 : p_1 = p_2$   $H_A : p_1 > p_2$   $\alpha = 0.05$   $\delta = 0.62$   $SE = 0.0346$  power = ?

*Step 1:* Find  $\hat{p}_1 - \hat{p}_2$  such that we reject  $H_0$ .

$$P(Z > z) < 0.05 \Rightarrow z > 1.644$$

$$\frac{(\hat{p}_1 - \hat{p}_2) - 0}{0.0346} > 1.644$$

$$\hat{p}_1 - \hat{p}_2 > 0.0569$$

What is the power of our hypothesis test to detect a difference of 0.62?

*Step 0:*  $H_0 : p_1 = p_2$   $H_A : p_1 > p_2$   $\alpha = 0.05$   $\delta = 0.62$   $SE = 0.0346$  power = ?

*Step 1:* Find  $\hat{p}_1 - \hat{p}_2$  such that we reject  $H_0$ .

$$P(Z > z) < 0.05 \Rightarrow z > 1.644$$

$$\frac{(\hat{p}_1 - \hat{p}_2) - 0}{0.0346} > 1.644$$

$$\hat{p}_1 - \hat{p}_2 > 0.0569$$

*Step 2:* Assume  $p_1 - p_2 = 0 + \delta = 0.62$  - we no longer assume independence, must use  $SE = 0.0273$  from the CI instead.

$$\begin{aligned} &P(\hat{p}_1 - \hat{p}_2 > 0.0569) \\ &= P\left(Z > \frac{0.0569 - 0.62}{0.0273}\right) \\ &= P(Z > -20) \end{aligned}$$

## Recap

---

## Recap - inference for one proportion

- Population parameter:  $p$ , point estimate:  $\hat{p}$

## Recap - inference for one proportion

- Population parameter:  $p$ , point estimate:  $\hat{p}$
- Conditions:
  - independence
    - random sample and 10% condition
  - at least 10 successes and failures
    - *observed* for CI
    - *expected* for HT



## Recap - inference for one proportion

- Population parameter:  $p$ , point estimate:  $\hat{p}$
- Conditions:
  - independence
    - random sample and 10% condition
  - at least 10 successes and failures
    - *observed* for CI
    - *expected* for HT
- Standard error:  $SE = \sqrt{\frac{p(1-p)}{n}}$ 
  - for CI: use  $\hat{p}$
  - for HT: use  $p_0$
  - for Power:
    - Step 1 - use  $p_0$
    - Step 2 - use  $p_A = p_0 + \delta$

## Recap - comparing two proportions

- Population parameter:  $(p_1 - p_2)$ , point estimate:  $(\hat{p}_1 - \hat{p}_2)$

## Recap - comparing two proportions

- Population parameter:  $(p_1 - p_2)$ , point estimate:  $(\hat{p}_1 - \hat{p}_2)$
- Conditions:

## Recap - comparing two proportions

- Population parameter:  $(p_1 - p_2)$ , point estimate:  $(\hat{p}_1 - \hat{p}_2)$
- Conditions:
  - independence within groups
    - random sample and 10% condition met for both groups
  - independence between groups
  - at least 10 successes and failures in each group
    - *observed* for CI
    - *expected* for HT

## Recap - comparing two proportions

- Population parameter:  $(p_1 - p_2)$ , point estimate:  $(\hat{p}_1 - \hat{p}_2)$
- Conditions:
  - independence within groups
    - random sample and 10% condition met for both groups
  - independence between groups
  - at least 10 successes and failures in each group
    - *observed* for CI
    - *expected* for HT
- $SE = \sqrt{\frac{p_1(1-p_1)}{n_1} + \frac{p_2(1-p_2)}{n_2}}$ 
  - for CI: use  $\hat{p}_1$  and  $\hat{p}_2$
  - for HT:
    - when  $H_0 : p_1 = p_2$ : use  $\hat{p}_{pool} = \frac{\#SUC_1 + \#SUC_2}{n_1 + n_2}$
    - when  $H_0 : p_1 - p_2 = (\text{some value other than } 0)$ : use  $\hat{p}_1$  and  $\hat{p}_2$ 
      - this is pretty rare
  - for Power:
    - Step 1 - use  $\hat{p}_{pool}$
    - Step 2 - use  $\hat{p}_1$  and  $\hat{p}_2$

## Reference - standard error calculations

	one sample	two samples
mean	$SE = \frac{\sigma}{\sqrt{n}}$	$SE = \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$
proportion	$SE = \sqrt{\frac{p(1-p)}{n}}$	$SE = \sqrt{\frac{p_1(1-p_1)}{n_1} + \frac{p_2(1-p_2)}{n_2}}$

## Reference - standard error calculations

	one sample	two samples
mean	$SE = \frac{\sigma}{\sqrt{n}}$	$SE = \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$
proportion	$SE = \sqrt{\frac{p(1-p)}{n}}$	$SE = \sqrt{\frac{p_1(1-p_1)}{n_1} + \frac{p_2(1-p_2)}{n_2}}$

- When working with means, it's very rare that  $\sigma$  is known, so we usually use  $s$  as an approximation.

## Reference - standard error calculations

	one sample	two samples
mean	$SE = \frac{\sigma}{\sqrt{n}}$	$SE = \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$
proportion	$SE = \sqrt{\frac{p(1-p)}{n}}$	$SE = \sqrt{\frac{p_1(1-p_1)}{n_1} + \frac{p_2(1-p_2)}{n_2}}$

- When working with means, it's very rare that  $\sigma$  is known, so we usually use  $s$  as an approximation.
- When working with proportions, we will not know  $p$  therefore
  - if doing a hypothesis test,  $p$  comes from the null hypothesis
  - if constructing a confidence interval, use  $\hat{p}$  instead