Midterm Examination #2

Mth 135 = Sta 104

Friday, 2001 November 16, 10:30 – 11:20 am

Please place all your books and notes on the floor. You may use a single one-sided sheet of prepared notes, if you wish, but you may not share materials. You may use a calculator but not a laptop, pda, or cell phone. If you find a question confusing please **ask me**, but do not talk to each other.

You must **show** your **work** to get partial credit. Unsupported answers are not acceptable, even if they are correct. Please give all numerical answers as fractions **in lowest terms** (simplify!) or as decimals correct to **four places**. You should spend about 10 minutes on each problem. It is to your advantage to write your solutions as clearly as possible, since I cannot give credit for solutions I do not understand. Good luck.

Cheating on exams is a breach of trust with classmates and faculty, and will not be tolerated. After completing the exam please acknowledge the Duke Honor Code:

I have neither given nor received any unauthorized aid on this exam.

Signature: ______________________

Print Name: ______________________

1.	/20
2.	/20
3.	/20
4.	/20
5.	/20
Total:	/100

Problem 1: Let Y have a continuous probability distribution with density function (pdf) $f_Y(y) = 1/2$, $0 < y < 2$; then let X have a conditional distribution (given Y) that is uniform on the interval from 0 to Y.

a) (5) Give the joint density function for X and Y; be sure to give your answer correctly for all x and all y:

$f(x, y) =$ ______________

b) (6) Find the marginal density functions for X and Y:

$f_X(x) =$ ______________ $f_Y(y) =$ ______________

c) (9) Find the indicated expectations. (Hint: Condition)

$\mathsf{E}[X\,Y] =$ ______________

$\mathsf{E}[X] =$ ______________

$\mathsf{E}[10 - 25\,Y] =$ ______________.

Problem 2: If X is a random variable with mean $\mathsf{E}[X] = 0$ and variance $\mathsf{E}[X^2] = 16$, and if Y is a random variable with mean $\mathsf{E}[Y = 0]$ and variance $\mathsf{E}[Y^2] = 9$, and if $\mathsf{E}[XY] = 3$, find:

a) (4) The mean and variance of $Z = 2X + 3$:

$\mathsf{E}[Z] =$ ____________________ $\mathsf{Var}[Z] =$ ____________________

b) (6) The mean and variance of $X - Y$:

$\mathsf{E}[X-Y] =$ ____________________ $\mathsf{Var}[X-Y] =$ ____________________

c) (5) The **correlation** (not *covariance*) between X and Y:

$\mathsf{Cor}(X, Y) =$ ____________________

d) (5) If X and Y are jointly normal, find the conditional expectation:

$\mathsf{E}[Y|X = 2] =$ ____________________

Problem 3: The random variables X and Y are independent with the same mean $\mu = 1$ and the same variance $\sigma^2 = 4$. Find:

a) (4) $\mathsf{E}[X - 2Y] =$ ____________

b) (4) $\mathsf{Var}[X - 2Y] =$ ____________

c) (4) $\mathsf{E}[X\,Y] =$ ____________

d) (4) $\mathsf{Var}[X\,Y] =$ ____________

e) (4) $\mathsf{P}\big[|X - Y| > 10\big] \le$ ____________ (find a bound)

Problem 4: Each of the random variables S and T has the same marginal distribution, with density function

$$f(u) = \begin{cases} u/200 & 0 < u < 10 \\ 0 & \text{otherwise} \end{cases}$$

a) (5) Find the probability density function for $X \equiv \sqrt{S}$.

$f_X(x) =$ ____________

b) (5) Find the indicated expectation:

$\mathsf{E}[e^S/S] =$ ________________

c) (5) Do you have enough information to find the indicated mean? If so, do it; if not, tell what's missing:

$\mathsf{E}[S - T] =$ ____________

d) (5) Do you have enough information to find the indicated variance? If so, do it; if not, tell what's missing:

$\mathsf{Var}[S - T] =$ ____________

Problem 5: For each $1 \leq j \leq 100$, the independent random variables X_j take on only the possible values zero, one, and two with probabilities $1/4$, $1/2$, and $1/4$, respectively. The random variable $S = X_1 + ... + X_{100}$ is their sum.

a) (6) Find the mean and variance of X_7:

$\mathsf{E}[X_7] =$ ______________ $\mathsf{Var}[X_7] =$ ______________

b) (6) Find the mean and variance of S:

$\mathsf{E}[S] =$ ______________ $\mathsf{Var}[S] =$ ______________

c) (8) Find the approximate probability:

$\mathsf{P}[S > 104] \approx$ ______________

d) (+2) Do you know the *exact* probability distribution of S? How?

$\mathsf{P}[S = s] =$ ______________

Extra worksheet, if needed:

$$\Phi(x) = \int_{-\infty}^{x} \frac{1}{\sqrt{2\pi}}\, e^{-z^2/2}\, dz:$$



Table 5.1 Area $\Phi(x)$ under the Standard Normal Curve to the left of x.

x	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998

$\Phi(0.6745) = 0.75$ $\Phi(1.6449) = 0.95$ $\Phi(2.3263) = 0.99$ $\Phi(3.0902) = 0.999$

$\Phi(1.2816) = 0.90$ $\Phi(1.9600) = 0.975$ $\Phi(2.5758) = 0.995$ $\Phi(3.2905) = 0.9995$

Name	**Notation**	**pdf/pmf**	**Range**	**Mean μ**	**Variance σ^2**	
Beta	$\mathsf{Be}(\alpha,\beta)$	$f(x)=\frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)}x^{\alpha-1}(1-x)^{\beta-1}$	$x\in(0,1)$	$\frac{\alpha}{\alpha+\beta}$	$\frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$	
Binomial	$\mathsf{Bi}(n,p)$	$f(x)=\binom{n}{x}p^x q^{(n-x)}$	$x\in 0,\cdots,n$	$n\,p$	$n\,p\,q$	$(q=1-p)$
Exponential	$\mathsf{Ex}(\lambda)$	$f(x)=\lambda\,e^{-\lambda x}$	$x\in\mathbb{R}_+$	$1/\lambda$	$1/\lambda^2$	
Gamma	$\mathsf{Ga}(\alpha,\lambda)$	$f(x)=\frac{\lambda^\alpha}{\Gamma(\alpha)}x^{\alpha-1}\,e^{-\lambda x}$	$x\in\mathbb{R}_+$	α/λ	α/λ^2	
Geometric	$\mathsf{Ge}(p)$	$f(x)=p\,q^x$	$x\in\mathbb{Z}_+$	q/p	q/p^2	$(q=1-p)$
		$f(x)=p\,q^{y-1}$	$y\in\{1,...\}$	$1/p$	q/p^2	$(y=x+1)$
HyperGeo.	$\mathsf{HG}(n,A,B)$	$f(x)=\frac{\binom{A}{x}\binom{B}{n-x}}{\binom{A+B}{n}}$	$x\in 0,\cdots,n$	$n\,P$	$n\,P\,(1-P)\frac{N-n}{N-1}$	$(P=\frac{A}{A+B})$
Logistic	$\mathsf{Lo}(\mu,\beta)$	$f(x)=\frac{e^{-(x-\mu)/\beta}}{\beta[1+e^{-(x-\mu)/\beta}]^2}$	$x\in\mathbb{R}$	μ	$\pi^2\beta^2/3$	
Log Normal	$\mathsf{LN}(\mu,\sigma^2)$	$f(x)=\frac{1}{x\sqrt{2\pi\sigma^2}}e^{-(\log x-\mu)^2/2\sigma^2}$	$x\in\mathbb{R}_+$	$e^{\mu+\sigma^2/2}$	$e^{2\mu+\sigma^2}\left(1-e^{\sigma^2}\right)$	
Neg. Binom.	$\mathsf{NB}(\alpha,p)$	$f(x)=\binom{x+\alpha-1}{x}p^\alpha\,q^x$	$x\in\mathbb{Z}_+$	$\alpha q/p$	$\alpha q/p^2$	$(q=1-p)$
		$f(x)=\binom{y-1}{y-\alpha}p^\alpha\,q^{y-\alpha}$	$y\in\{\alpha,...\}$	α/p	$\alpha q/p^2$	$(y=x+\alpha)$
Normal	$\mathsf{No}(\mu,\sigma^2)$	$f(x)=\frac{1}{\sqrt{2\pi\sigma^2}}e^{-(x-\mu)^2/2\sigma^2}$	$x\in\mathbb{R}$	μ	σ^2	
Pareto	$\mathsf{Pa}(\alpha,\beta)$	$f(x)=\beta\,\alpha^\beta/x^{\beta+1}$	$x\in(\alpha,\infty)$	$\frac{\alpha\,\beta}{\beta-1}$	$\frac{\alpha^2\beta}{(\beta-1)^2(\beta-2)}$	
Poisson	$\mathsf{Po}(\lambda)$	$f(x)=\frac{\lambda^x}{x!}e^{-\lambda}$	$x\in\mathbb{Z}_+$	λ	λ	
Snedecor F	$F(\nu_1,\nu_2)$	$f(x)=\frac{\Gamma(\frac{\nu_1+\nu_2}{2})(\nu_1/\nu_2)^{\nu_1/2}}{\Gamma(\frac{\nu_1}{2})\Gamma(\frac{\nu_2}{2})}\times$	$x\in\mathbb{R}_+$	$\frac{\nu_2}{\nu_2-2}$	$\left(\frac{\nu_2}{\nu_2-2}\right)^2\frac{2(\nu_1+\nu_2-2)}{\nu_1(\nu_2-4)}$	
		$x^{\frac{\nu_1-2}{2}}\left[1+\frac{\nu_1}{\nu_2}x\right]^{-\frac{\nu_1+\nu_2}{2}}$				
Student t	$t(\nu)$	$f(x)=\frac{\Gamma(\frac{\nu+1}{2})}{\Gamma(\frac{\nu}{2})\sqrt{\pi\nu}}[1+x^2/\nu]^{-(\nu+1)/2}$	$x\in\mathbb{R}$	0	$\nu/(\nu-2)$	
Uniform	$\mathsf{Un}(a,b)$	$f(x)=\frac{1}{b-a}$	$x\in(a,b)$	$\frac{a+b}{2}$	$\frac{(b-a)^2}{12}$	
Weibull	$\mathsf{We}(\alpha,\beta,\gamma)$	$f(x)=\frac{\alpha(x-\gamma)^{\alpha-1}}{\beta^\alpha}e^{-[(x-\gamma)/\beta]^\alpha}$	$x\in(\gamma,\infty)$	$\gamma+\beta\Gamma(1+\alpha^{-1})$		