Midterm Examination # 2_a

Mth 135 = Sta 104

Thursday, 2003 November 20, 9:10 – 10:25 am

Please place all your books and notes on the floor. You may use a calculator and a single one-sided sheet of prepared notes, if you wish, but you may not use any other device and may not share materials. If you find a question confusing **please ask me**, but do not talk to each other.

You must **show** your **work** to get partial credit. Unsupported answers are not acceptable, even if they are correct. Please give all numerical answers as fractions **in lowest terms** (simplify!) or as decimals correct to **four places**. You should spend about 10 minutes on each problem. It is to your advantage to write your solutions as clearly as possible, since I cannot give credit for solutions I do not understand. Good luck.

Cheating on exams is a breach of trust with classmates and faculty, and will not be tolerated. After completing the exam please acknowledge that you have adhered to the Duke Community Standard:

- I will not lie, cheat, or steal in my academic endeavors, nor will I accept the actions of those who do.
- I will conduct myself responsibly and honorably in all my activities as a Duke student.

Signature: ______________________

Print Name: ______________________

1.	/20
2.	/20
3.	/20
4.	/20
5.	/20
Total:	/100

Problem 1: Let Y have a continuous probability distribution with density function (pdf) $f_Y(y) = 2y$, $0 < y < 1$; then let X have a conditional distribution (given Y) that is uniform on the interval from 0 to Y.

a) (5) Give the joint density function for X and Y; be sure to give your answer correctly for all x and all y:

$$f(x, y) = \begin{cases} \\ \\ \end{cases}$$

b) (6) Find the marginal density functions for X and Y:

$$f_X(x) = \begin{cases} \\ \\ \end{cases} \qquad f_Y(y) = \begin{cases} \\ \\ \end{cases}$$

c) (6) Find the conditional density functions for X and Y:

$$f_{X|Y}(x \mid y) = \begin{cases} \\ \\ \end{cases} \qquad f_{Y|X}(y \mid x) = \begin{cases} \\ \\ \end{cases}$$

d) (3) Find the expectation:

$\mathsf{E}[e^X \mid Y = 1/2] =$ ______________

Problem 2: The random variables X and Y are independent with the same mean $\mu = 1$ and the same variance $\sigma^2 = 4$. If possible, find:

a) (4) $\mathsf{E}[X - 2Y] =$ ______________

b) (4) $\mathsf{E}[X^2] =$ ______________

c) (4) $\mathsf{E}[X\,Y] =$ ______________

d) (4) $\mathsf{E}[(X - Y)^2] =$ ______________

e) (4) $\mathsf{Var}[X + 2\,Y] =$ ______________

Problem 3: If X is a random variable with mean $\mathsf{E}[X] = 0$ and variance $\mathsf{E}[X^2] = 16$, and if Y is a random variable with mean $\mathsf{E}[Y] = 0$ and variance $\mathsf{E}[Y^2] = 9$, and if $\mathsf{E}[XY] = 3$, find:

a) (4) The mean and variance of $Z = 2X + 3$:

$\mathsf{E}[Z] =$ ______________ $\mathsf{Var}[Z] =$ ______________

b) (6) The mean and variance of $X-Y$:

$\mathsf{E}[X-Y] =$ ______________ $\mathsf{Var}[X-Y] =$ ______________

c) (5) The **correlation** (not *covariance*) between X and Y:

$\mathsf{Cor}(X, Y) =$ ______________

d) (5) If X and Y are jointly normal, find the conditional expectation:

$\mathsf{E}[Y|X = 2] =$ ______________

Problem 4: Each of the random variables S and T has the same marginal distribution, with density function

$$f(u) = \begin{cases} 3\,u^{-4} & u > 1 \\ 0 & \text{otherwise} \end{cases}$$

a) (5) Find the probability density function for $X \equiv \sqrt{S}$.

$$f_X(x) = \begin{cases} & \\ & \end{cases}$$

b) (5) Find the indicated expectation:

$\mathsf{E}[1/S] =$ ____________

c) (5) Do you have enough information to find the indicated mean? If so, do it; if not, tell what's missing:

$\mathsf{E}[2S - T] =$ ____________

d) (5) Do you have enough information to find the indicated variance? If so, do it; if not, tell what's missing:

$\mathsf{Var}[2S - T] =$ ____________

Problem 5: For each $1 \le j \le 200$, the independent random variables X_j take on only the possible values -2, zero, and $+3$, with probabilities $1/4$, $1/4$, and $1/2$, respectively.

a) (6) Find the mean and variance of X_8:

$\mathsf{E}[X_8] =$ ______________ $\mathsf{Var}[X_8] =$ ______________

b) (6) Find the mean and variance of the sum $S = X_1 + ... + X_{200}$:

$\mathsf{E}[S] =$ ______________ $\mathsf{Var}[S] =$ ______________

c) (8) Find (to 4 decimal places) the approximate probability:

$\mathsf{P}[S > 215] \approx$ ______________

Extra worksheet, if needed:

$$\Phi(x) = \int_{-\infty}^{x} \frac{1}{\sqrt{2\pi}} e^{-z^2/2}\, dz:$$



Table 5.1 Area $\Phi(x)$ under the Standard Normal Curve to the left of x.

x	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998

$\Phi(0.6745) = 0.75$ $\Phi(1.6449) = 0.95$ $\Phi(2.3263) = 0.99$ $\Phi(3.0902) = 0.999$
$\Phi(1.2816) = 0.90$ $\Phi(1.9600) = 0.975$ $\Phi(2.5758) = 0.995$ $\Phi(3.2905) = 0.9995$

Name	Notation	pdf/pmf	Range	Mean μ	Variance σ^2	
Beta	$\mathsf{Be}(\alpha,\beta)$	$f(x)=\frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)}x^{\alpha-1}(1-x)^{\beta-1}$	$x\in(0,1)$	$\frac{\alpha}{\alpha+\beta}$	$\frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$	
Binomial	$\mathsf{Bi}(n,p)$	$f(x)=\binom{n}{x}p^x q^{(n-x)}$	$x\in 0,\cdots,n$	$n\,p$	$n\,p\,q$	$(q=1-p)$
Exponential	$\mathsf{Ex}(\lambda)$	$f(x)=\lambda\,e^{-\lambda x}$	$x\in\mathbb{R}_+$	$1/\lambda$	$1/\lambda^2$	
Gamma	$\mathsf{Ga}(\alpha,\lambda)$	$f(x)=\frac{\lambda^\alpha}{\Gamma(\alpha)}x^{\alpha-1}\,e^{-\lambda x}$	$x\in\mathbb{R}_+$	α/λ	α/λ^2	
Geometric	$\mathsf{Ge}(p)$	$f(x)=p\,q^x$	$x\in\mathbb{Z}_+$	q/p	q/p^2	$(q=1-p)$
		$f(x)=p\,q^{y-1}$	$y\in\{1,...\}$	$1/p$	q/p^2	$(y=x+1)$
HyperGeo.	$\mathsf{HG}(n,A,B)$	$f(x)=\frac{\binom{A}{x}\binom{B}{n-x}}{\binom{A+B}{n}}$	$x\in 0,\cdots,n$	$n\,P$	$n\,P\,(1-P)\frac{N-n}{N-1}$	$(P=\frac{A}{A+B})$
Logistic	$\mathsf{Lo}(\mu,\beta)$	$f(x)=\frac{e^{-(x-\mu)/\beta}}{\beta[1+e^{-(x-\mu)/\beta}]^2}$	$x\in\mathbb{R}$	μ	$\pi^2\beta^2/3$	
Log Normal	$\mathsf{LN}(\mu,\sigma^2)$	$f(x)=\frac{1}{x\sqrt{2\pi\sigma^2}}e^{-(\log x-\mu)^2/2\sigma^2}$	$x\in\mathbb{R}_+$	$e^{\mu+\sigma^2/2}$	$e^{2\mu+\sigma^2}\left(1-e^{\sigma^2}\right)$	
Neg. Binom.	$\mathsf{NB}(\alpha,p)$	$f(x)=\binom{x+\alpha-1}{x}p^\alpha\,q^x$	$x\in\mathbb{Z}_+$	$\alpha q/p$	$\alpha q/p^2$	$(q=1-p)$
		$f(x)=\binom{y-1}{y-\alpha}p^\alpha\,q^{y-\alpha}$	$y\in\{\alpha,...\}$	α/p	$\alpha q/p^2$	$(y=x+\alpha)$
Normal	$\mathsf{No}(\mu,\sigma^2)$	$f(x)=\frac{1}{\sqrt{2\pi\sigma^2}}e^{-(x-\mu)^2/2\sigma^2}$	$x\in\mathbb{R}$	μ	σ^2	
Pareto	$\mathsf{Pa}(\alpha,\beta)$	$f(x)=\beta\,\alpha^\beta/x^{\beta+1}$	$x\in(\alpha,\infty)$	$\frac{\alpha\,\beta}{\beta-1}$	$\frac{\alpha^2\beta}{(\beta-1)^2(\beta-2)}$	
Poisson	$\mathsf{Po}(\lambda)$	$f(x)=\frac{\lambda^x}{x!}e^{-\lambda}$	$x\in\mathbb{Z}_+$	λ	λ	
Snedecor F	$F(\nu_1,\nu_2)$	$f(x)=\frac{\Gamma(\frac{\nu_1+\nu_2}{2})(\nu_1/\nu_2)^{\nu_1/2}}{\Gamma(\frac{\nu_1}{2})\Gamma(\frac{\nu_2}{2})}\times$ $x^{\frac{\nu_1-2}{2}}\left[1+\frac{\nu_1}{\nu_2}\,x\right]^{-\frac{\nu_1+\nu_2}{2}}$	$x\in\mathbb{R}_+$	$\frac{\nu_2}{\nu_2-2}$	$\left(\frac{\nu_2}{\nu_2-2}\right)^2\frac{2(\nu_1+\nu_2-2)}{\nu_1(\nu_2-4)}$	
Student t	$t(\nu)$	$f(x)=\frac{\Gamma(\frac{\nu+1}{2})}{\Gamma(\frac{\nu}{2})\sqrt{\pi\nu}}[1+x^2/\nu]^{-(\nu+1)/2}$	$x\in\mathbb{R}$	0	$\nu/(\nu-2)$	
Uniform	$\mathsf{Un}(a,b)$	$f(x)=\frac{1}{b-a}$	$x\in(a,b)$	$\frac{a+b}{2}$	$\frac{(b-a)^2}{12}$	
Weibull	$\mathsf{We}(\alpha,\beta,\gamma)$	$f(x)=\frac{\alpha(x-\gamma)^{\alpha-1}}{\beta^\alpha}e^{-[(x-\gamma)/\beta]^\alpha}$	$x\in(\gamma,\infty)$	$\gamma+\beta\Gamma(1+\alpha^{-1})$		