

STA 711: Note from Nov 13

Let $\{X_n\}$ be independent random variables for $n \in \mathbb{N}$ with

$$\mathbb{P}[X_n = x] = \begin{cases} n^{-1} & x = 1 \\ 1 - n^{-1} & x = 0. \end{cases}$$

In what sense(s) does $T_m := \sum_{1 \leq n \leq m} n^{-1} X_n$ converge to a finite random variable T as $m \rightarrow \infty$?

Convergence in L_1 to $T := \sum_{1 \leq n < \infty} n^{-1} X_n$ is easy to show, either using the monotone convergence theorem and the calculation

$$\mathbb{E}[T] = \sum_{n=1}^{\infty} \mathbb{E}[X_n/n] = \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6} < \infty$$

or the explicit bound

$$\mathbb{E}|T - T_m| = \sum_{m+1}^{\infty} \frac{1}{n^2} < \int_m^{\infty} \frac{1}{x^2} dx = \frac{1}{m} \rightarrow 0.$$

It follows immediately that T_m converges almost-surely, because T is infinite on the set

$$\mathcal{N} = \{\omega : T_m(\omega) \text{ does not converge}\}.$$

Since $T \in L_1$, necessarily $\mathbb{P}[\mathcal{N}] = 0$, and $T_m \rightarrow T$ *a.s.* (and so also *pr.*).

For any $1 \leq p < \infty$, Minkowski's inequality and the calculation $\|X_n\|_p = (1/n)^{1/p} = n^{-1/p}$ (so $\|X_n/n\|_p = n^{-1-1/p}$) imply

$$\begin{aligned} \|T - T_m\|_p &= \left\| \sum_{m+1}^{\infty} X_n/n \right\|_p \leq \sum_{m+1}^{\infty} \|X_n/n\|_p \\ &= \sum_{m+1}^{\infty} n^{-1-1/p} < \int_m^{\infty} x^{-1-1/p} dx = pm^{-1/p} \rightarrow 0, \end{aligned}$$

so also $T_m \rightarrow T$ in L_p for all $1 \leq p < \infty$.

It doesn't converge in L_∞ , though, because for any $B < \infty$ and any N large enough that $\sum_{m < n \leq N} \frac{1}{n} > B$ (always possible since the harmonic series diverges),

$$\begin{aligned} \mathbb{P}[|T - T_m| > B] &\geq \mathbb{P}[X_n = 1 \text{ for } m < n \leq N] \\ &\geq \prod_{m < n \leq N} \frac{1}{n} = \frac{m!}{N!} > 0. \end{aligned}$$

The smallest possible choice will be approximately $N \approx me^B$.