

Sta 711: Homework 9

Uniform Integrability

1. **True or false?** Answer whether each of the following statements is true or false. If true, answer why; if false, give a simple counter example.
 - (a) If $\{X_n, n \in \mathbb{N}\}$ is a uniformly integrable (UI) collection of random variables, then X_n is uniformly bounded in L_1 .
 - (b) Define a sequence $\{X_n\}$ of random variables on the unit interval with Lebesgue measure, (Ω, \mathcal{F}, P) with $\Omega = (0, 1]$, $\mathcal{F} = \mathcal{B}$, and $P = \lambda$, by $X_n := \sqrt{n} \mathbf{1}_{(0, \frac{1}{n}]}$. Then $\{X_n\}$ is UI.
 - (c) Let $\{X_n\}$ be a sequence of random variables for which $e^{|X_n|}$ is uniformly bounded in L_1 , *i.e.*, satisfies $\mathbb{E}e^{|X_n|} \leq B$ for some $B < \infty$ and all n . Then $\{X_n\}$ is UI.
 - (d) Let $\{X_n\}$ be a sequence of random variables that is uniformly bounded in L_1 , *i.e.*, satisfies $\mathbb{E}|X_n| \leq B$ for some $B < \infty$ and all n . Then $\{X_n\}$ is UI.

Characteristic Functions

2. Let X be a random variable, and define

$$\phi_X(\theta) := \mathbb{E}(e^{i\theta X}), \quad \theta \in \mathbb{R}$$

Show that $\phi_X(\theta)$ is uniformly continuous in \mathbb{R} .

3. Find the characteristic functions of the following random variables:

- (a) $W := c^1$ (The superscripts in (a)–(c) are footnote indicators, not exponents)
- (b) $X \sim \text{Un}(a, b)^2$
- (c) $Y \sim \text{Ga}(\alpha, \lambda)^3$
- (d) $Z_n = (Y_1 + Y_2 + \cdots + Y_n)/n, \quad Y_j \stackrel{\text{iid}}{\sim} \text{Ga}(\alpha, \lambda)$

What is the distribution of Z_n ? What happens as $n \rightarrow \infty$?

4. The distribution of a random variable X is called *infinitely divisible* if, for every $n \in \mathbb{N}$, there exist n iid random variables $\{Y_i\}$ such that X has the same distribution as $\sum_{i=1}^n Y_i$. Use characteristic functions to show that if $X \sim \text{Po}(\lambda)$, then X is infinitely divisible.⁴

¹A constant random variable with value $c \in \mathbb{R}$

²Uniform, on the interval $(a, b) \subset \mathbb{R}$

³Gamma, with rate parameterization— with pdf $f(y | \lambda) = \lambda^\alpha y^{\alpha-1} e^{-\lambda y} / \Gamma(\alpha)$, $y > 0$.

⁴Hint: If $\{Y_i\}$ are independent with sum $Y_+ := \sum Y_i$, then $\phi_{Y_+}(\theta) = \prod \phi_{Y_i}(\theta)$ for all $\theta \in \mathbb{R}$.

5. Suppose $\{A_n, n \in \mathbb{N}\}$ are independent events satisfying $\mathbf{P}(A_n) < 1, \forall n \in \mathbb{N}$. Show that $\mathbf{P}(\bigcup_{n=1}^{\infty} A_n) = 1$ if and only if $\mathbf{P}(A_n \text{ i.o.}) = 1$ (“i.o.” means “infinitely often”, so the question concerns $\limsup A_n$). Give an example to show that the condition $\mathbf{P}(A_n) < 1$ cannot be dropped.
6. Let $\{A_n\}$ be a sequence of events with $\mathbf{P}(A_n) \rightarrow 1$ as $n \rightarrow \infty$. Prove that there exists a subsequence $\{n_k\}$ tending to infinity such that $\mathbf{P}(\bigcap_k A_{n_k}) > 0$.
7. Let A_n be a sequence of events that all satisfy $\mathbf{P}(A_n) \geq \epsilon$ for some $\epsilon > 0$. Does there necessarily exist a subsequence $\{n_k \rightarrow \infty\}$ with $\mathbf{P}(\bigcap_k A_{n_k}) > 0$? Why or why not?
8. Let $\{X_n\}$ be non-negative iid random variables, with tail σ -field

$$\mathcal{T} := \bigcap_{n \in \mathbb{N}} \mathcal{F}'_n, \quad \mathcal{F}'_n := \sigma\{X_m : m > n\}$$

Is the event

$$\begin{aligned} E &= \{\text{There exists } \epsilon > 0 \text{ such that } X_n > n\epsilon \text{ for infinitely-many } n\} \\ &= \bigcup_{\epsilon > 0} \bigcap_{n \geq 1} \bigcup_{m \geq n} \{\omega : X_m(\omega) > m\epsilon\} \end{aligned}$$

in \mathcal{T} ? Prove or disprove it.

Express the probability $\mathbf{P}[E]$ in terms of the random variables' common distribution—for example, using their common CDF $F(x) := \mathbf{P}[X_n \leq x]$ or moments $\mathbf{E}[|X_n|^p]$ for some $p > 0$.