

## Sta 711: Homework 6

### Independence

1. Let  $\{B_i\}$  be independent events. For  $n \in \mathbb{N}$  show that

$$\mathbb{P}\left(\bigcup_{i=1}^n B_i\right) = 1 - \prod_{i=1}^n [1 - \mathbb{P}(B_i)] \geq 1 - \exp\left\{-\sum_{i=1}^n \mathbb{P}(B_i)\right\}$$

and conclude that  $\mathbb{P}[\bigcup_{i=1}^{\infty} B_i] = 1$  if each  $\mathbb{P}[B_i] \geq \epsilon$  for some  $\epsilon > 0$ . Show that this conclusion would be false without the assumption of independence.

2. If  $\{A_n, n \in \mathbb{N}\}$  is a sequence of events such that  $\mathbb{P}[A_n] = 1/3$  for each  $n$  and

$$(\forall n \neq m \in \mathbb{N}) \quad \mathbb{P}(A_n \cap A_m) = \mathbb{P}(A_n)\mathbb{P}(A_m),$$

does it follow that the events  $\{A_n\}$  are independent? Give a proof or counter-example. Note  $1/3 \neq 1/2$ .

3. Show that a random variable  $Y$  is independent of itself if and only if, for some constant  $c \in \mathbb{R}$ ,  $\mathbb{P}[Y = c] = 1$ .  
Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be Borel measurable, and  $X$  a non-constant random variable. Can  $Y := f(X)$  and  $X$  be independent? Explain your answer.
4. Give an example to show that an event  $A \in \mathcal{F}$  may be independent of each  $B$  in some collection  $\mathcal{C} \subset \mathcal{F}$  of events, but *not* independent of  $\sigma(\mathcal{C})$ . Prove this is impossible if  $\mathcal{C}$  is a  $\pi$ -system (*i.e.*, in that case  $A$  must be independent of  $\sigma(\mathcal{C})$ ).
5. Give a simple example to show that two random variables on the same space  $(\Omega, \mathcal{F})$  may be independent according to one probability measure  $\mathbb{P}_1$  but dependent with respect to another  $\mathbb{P}_2$ .

### Fubini's Theorem

6. Let  $X \geq 0$  be a positive random variable and  $\alpha > 0$ . Show that

$$\mathbb{E}(X^\alpha) = \alpha \int_0^\infty t^{\alpha-1} \mathbb{P}(X > t) dt.$$

Note that the distribution  $\mu(dx)$  of  $X$  need not be absolutely continuous (so  $X$  may not have a pdf). Where did you use Fubini's theorem?

7. Define measure spaces  $(\Omega_i, \mathcal{F}_i, \mu_i)$ , for  $i = 1, 2$  as follows. Let each  $\Omega_i := (0, 1]$ , the unit interval, with  $\sigma$ -algebras

$$\mathcal{F}_1 = \mathcal{B} = \text{Borel sets of } (0,1] \quad \mathcal{F}_2 = 2^\Omega = \text{All subsets of } (0,1],$$

and let  $\mu_1 = \lambda$  be Lebesgue measure and  $\mu_2$  counting measure— so  $\mu_1(A)$  is the length of any Borel set  $A \in \mathcal{F}_1$  and  $\mu_2(B)$  is the cardinality of  $B \subset (0, 1]$ . Define

$$f(x, y) := \mathbf{1}_{x=y}(x, y)$$

Set

$$I_1 := \int_{\Omega_1} \left[ \int_{\Omega_2} f(x, y) \mu_2(dy) \right] \mu_1(dx) \quad I_2 := \int_{\Omega_2} \left[ \int_{\Omega_1} f(x, y) \mu_1(dx) \right] \mu_2(dy)$$

Compute  $I_1$  and  $I_2$ . Is  $I_1 = I_2$ ? Are the measures  $\mu_1$  and  $\mu_2$   $\sigma$ -finite? Why doesn't Fubini's theorem hold here?

8. This problem is a probabilistic version of the familiar integration-by-parts formula from calculus. Suppose  $F$  and  $G$  are two distribution functions with no common points of discontinuity on an interval  $(a, b]$ . Show that

$$\int_{(a,b]} G(x)F(dx) = F(b)G(b) - F(a)G(a) - \int_{(a,b]} F(x)G(dx)$$

where “ $G(dx)$ ” denotes the measure on  $(\mathbb{R}, \mathcal{B})$  with DF  $G(x)$ . Show that the formula fails if  $F$  and  $G$  have common discontinuities.

### Zero-One Laws

9. Let  $\{X_n\}$  be a sequence of Bernoulli random variables with

$$\mathbb{P}(X_n = 1) = n^{-p} \quad \mathbb{P}(X_n = 0) = 1 - n^{-p}$$

for some  $p > 0$ . For  $p = 2$  show that the partial sum

$$S_n := \sum_{k=1}^n X_k$$

converges almost-surely, whether or not the  $\{X_n\}$  are independent. If the  $\{X_n\}$  are independent, for which  $p > 0$ , does  $S_n$  converge? Why?

10. Let  $\{X_n\}$  be an iid sequence of random variables with a non-degenerate distribution (*i.e.*, for some  $B \in \mathcal{B}$ ,  $0 < \mathbb{P}[X_n \in B] < 1$ ). Show that

$$\mathbb{P}[\omega : X_n(\omega) \text{ converges}] = 0$$

11. Use the Borel-Cantelli lemma to prove that for any sequence of real-valued random variables  $\{X_n\}$  (not necessarily independent or identically-distributed), there exist constants  $c_n \rightarrow \infty$  such that

$$\mathbb{P} \left( \lim_{n \rightarrow \infty} \frac{X_n}{c_n} = 0 \right) = 1.$$

Give a careful description of how you choose  $c_n$  (it will depend on the distributions of the  $X_n$ ). Find a suitable sequence  $\{c_n\}$  explicitly for an iid sequence  $\{X_n\} \stackrel{\text{iid}}{\sim} \text{Ex}(1)$  of unit-rate exponentially-distributed random variables to ensure that  $X_n/c_n \rightarrow 0$  almost surely.