

# Sta 711: Homework 5

## Convergence

1. Let  $X$  be a strictly positive random variable. Show that:

- (a)  $\lim_{n \rightarrow \infty} n \mathbb{E}(\frac{1}{X} \mathbf{1}_{[X > n]}) = 0$ .
- (b)  $\lim_{n \rightarrow \infty} n^{-1} \mathbb{E}(\frac{1}{X} \mathbf{1}_{[X > n^{-1}]}) = 0$ .

2. Let  $X \sim \text{Un}(0, 4]$  be uniformly distributed on the interval  $(0, 4]$ , and set  $Y := 1/X$  and  $Z := \log(4Y)$ . Suggestion: First find out what is the distribution of  $Z$ , by computing  $\mathbb{P}[Z > z]$  for  $z \in \mathbb{R}$ . Use  $\varphi(x) := |x|$  for the Markov inequality questions.

- (a) What bound does Markov's inequality give for  $\mathbb{P}[X > 3]$ ?
- (b) What bound does Chebychev's inequality give for  $\mathbb{P}[|X - 2| > 1]$ ?
- (c) What bound does Markov's inequality give for  $\mathbb{P}[Y > 1]$ ?
- (d) What bound does Markov's inequality give for  $\mathbb{P}[Z > 2]$ ?
- (e) What are the exact values of  $\mathbb{P}[X > 3]$ ,  $\mathbb{P}[|X - 2| > 1]$ ,  $\mathbb{P}[Y > 1]$ , and  $\mathbb{P}[Z > 2]$ ?

3. Let  $A$  and  $B$  be events in  $(\Omega, \mathcal{F}, \mathbb{P})$  with probabilities  $a = \mathbb{P}(A)$  and  $b = \mathbb{P}(B)$  respectively. Show that  $\mathbb{P}(A \cap B) \leq \sqrt{ab}$ .

4. Suppose  $\{X_n\}, X$  are real valued RVs defined on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  and that  $X_n(\omega) \rightarrow X(\omega)$  for all  $\omega \in \Omega$ . Show that for every  $\epsilon > 0$ , there is an event  $\Lambda_\epsilon$  with  $\mathbb{P}(\Lambda_\epsilon) < \epsilon$  and

$$\sup_{\omega \in \Lambda_\epsilon^c} |X(\omega) - X_n(\omega)| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Thus the convergence is uniform except on an arbitrarily small set. (For more on this result, called Egorov's Theorem, see page 89 of the text.)

5. For a random variable  $X$ ,  $1 < p < q < \infty$ , show<sup>1</sup> that

$$0 \leq \|X\|_1 \leq \|X\|_p \leq \|X\|_q \leq \|X\|_\infty$$

6. For  $1 < p < q < \infty$ , show that

$$L_\infty \subset L_q \subset L_p \subset L_1$$

where  $L_p := \{X : \|X\|_p < \infty\}$ .

---

<sup>1</sup>Hint: Jensen's inequality may help for some parts

7. The “Moment Generating Function” (MGF) of a real-valued random variable  $X$  (or of its distribution  $\mu(dx)$ ) is the extended real-valued function  $M_X(t) := \mathbf{E} \exp(tX) = \int_{\mathbb{R}} e^{tx} \mu(dx)$  of  $t \in \mathbb{R}$ . Show that a nonnegative random variable  $X \geq 0$  is in  $L_1$  if there exists some  $t > 0$  for which  $M_X(t) < \infty$ . Show that the converse may fail— *i.e.*, there exist  $X \geq 0$  in  $L_1$  for which  $M_X(t) = \infty$  for all  $t > 0$ .
8. Show that Minkowski’s Inequality fails for  $0 < p < 1$ — *i.e.*, find  $(\Omega, \mathcal{F}, \mathbf{P})$  and  $X, Y \in L_p(\Omega, \mathcal{F}, \mathbf{P})$  for which  $\|X + Y\|_p > \|X\|_p + \|Y\|_p$  for some  $0 < p < 1$ .