

Midterm Examination # 2

Mth 136 = Sta 114

Monday, April 16, 2009

2:50 – 4:05 pm

Version *b*

- This is a **closed book** exam— put your books on the floor.
- You may use a calculator and **one page** of your own notes.
- Do not share calculators or notes.
- **Show your work.** Neatness counts. Boxing answers helps.
- Numerical answers: **four significant digits** or simplified exact solution (e.g., fractions **in lowest terms**).
Simplify *all* answers for full credit.
- Extra worksheet and pdf & distribution tables are attached.

Cheating on exams is a breach of trust with classmates and faculty that will not be tolerated. After completing the exam please acknowledge that you have adhered to the Duke Community Standard:

- I will not lie, cheat, or steal in my academic endeavors;
- I will conduct myself honorably in all my endeavors; and
- I will act if the Standard is compromised.

Signature: _____

Print Name: _____

1.	/20
2.	/20
3.	/20
4.	/20
5.	/20
Total:	/100


Problem 1: Skyler thinks X has a standard exponential distribution with density function

$$f_0(x) = e^{-x} \mathbf{1}_{\{x>0\}},$$

while Terry thinks it has a lognormal distribution (so $Y \equiv \log X$ has a standard Normal distribution $Y \sim \text{No}(0, 1)$, $X = e^Y$) with density function

$$f_1(x) = \frac{1}{x\sqrt{2\pi}} e^{-(\log x)^2/2} \mathbf{1}_{\{x>0\}}.$$

When they disagree about things, Skyler is right about $2/3$ of the time. Here's a plot of the likelihood ratio against H_0 , $\Lambda(x) = f_1(x)/f_0(x)$, vs. x :



Problem 1 (cont):

- a) (6) Prove that the likelihood ratio increases monotonically on all of $0 < x < \infty$.

- b) (6) Based on the information above and the single observation $X = 3$, what is the (posterior) probability that Skyler was right? Give answer to four significant digits (or exact answer), and **simplify**:

$$P[\text{Skyler correct} \mid X = 3] =$$

Problem 1 (cont):

- c) (6) Find the P -value for likelihood ratio test of $H_0 : X \sim f_0(x)$ against alternative $H_1 : X \sim f_1(x)$ for the single observation $X = 3$ (as usual, exact or 4 sig digs:

$$P =$$

- c) (1) Would you Accept or Reject H_0 at level $\alpha = 0.05$?

- d) (1) Is the test in c) above UMP? Circle one: Y N
Why or why not?

Problem 2: The $n = 9$ independent random variables $\{X_i\}_{i \leq n}$ are a simple random sample from the Normal distribution $X_i \sim \mathbf{No}(\mu, \sigma^2)$, with variance σ^2 and mean μ .

- a) (7) With σ^2 unknown, describe the Rejection Region \mathcal{R} for a two-sided test of size $\alpha = 0.05$ of the hypothesis

$$H_0 : \mu = 17.5 \quad \text{vs.} \quad H_1 : \mu \neq 17.5$$

Reject H_0 if:

- b) (7) Same question, if variance of $\{X_i\}$ is known to be $\sigma^2 = 16$:

Reject H_0 if:

- c) (6) What is the *power* $\pi(\mu)$ of the test in part b) above, for $\mu = 20$?

$$\pi(20) = \underline{\hspace{2cm}}$$

Problem 3: Once again the $n = 9$ independent random variables $\{X_i\}_{i \leq n}$ are a simple random sample from the Normal distribution $X_i \sim \mathbf{No}(\mu, \sigma^2)$, with variance σ^2 and mean μ , but now we have some data. The values of a few statistics from this random sample are:

$$\begin{aligned} S(\mathbf{x}) = \sum_{i \leq n} X_i &= 180 & T(\mathbf{x}) = \min_{i \leq n} X_i &= 1 \\ V(\mathbf{x}) = \sum_{i \leq n} (X_i - \bar{X})^2 &= 72 & W(\mathbf{x}) = \text{Median}(\{X_i\}) &= 24 \end{aligned}$$

- a) (7) With σ^2 unknown, give the P -value for a two-sided test of the hypothesis

$$H_0 : \mu = 17.5 \quad \text{vs.} \quad H_1 : \mu \neq 17.5$$

$P =$

- b) (7) Same question, if variance of $\{X_i\}$ is known to be $\sigma^2 = 16$:

$P =$

- c) (6) Are these answers consistent with your answers to parts a) and b) of Problem 2? **Y N** Explain.

Problem 4: For some parameter value $0 < \theta < \infty$, the three random variables $\{X_i\}_{1 \leq i \leq 3}$ are uniformly distributed on the interval $[0, \theta]$. A few statistics from this random sample to consider are:

$$S(\mathbf{x}) = \sum X_i \quad T(\mathbf{x}) = \sum (X_i - \bar{X})^2 \quad V(\mathbf{x}) = \min X_i \quad W(\mathbf{x}) = \max X_i$$

- a) (6) If θ is known to be one of $\{\theta_0 = 2, \theta_1 = 3\}$, with prior probabilities $\xi_0 = 0.8$ and $\xi_1 = 0.2$ respectively, find the posterior probability indicated, as a function of one or more of S, T, V, W . Simplify!

$$\xi(\theta = \theta_0 \mid \mathbf{x}) =$$

- b) (6) If (instead) θ might be any number in $[1, \infty)$, with a Pareto prior distribution $\theta \sim \text{Pa}(\alpha, \epsilon)$ with density function

$$\xi(\theta) = \alpha \epsilon^\alpha / \theta^{\alpha+1} \mathbf{1}_{\{\theta \geq \epsilon\}}$$

with $\alpha = 1$ and $\epsilon = 1$, find the posterior density function for θ , correctly for all $\theta \in \mathbb{R}$. If possible, give your answer using one or more of S, T, V, W . Simplify!

$$\xi(\theta \mid \mathbf{x}) =$$

Problem 4 (cont):

- c) (8) Now suppose we observe data $\mathbf{x} = \{0.60, 1.75, 1.25\}$ and we wish to make a decision about the hypothesis $H_0 : \theta \leq 2$ with alternative $H_1 : \theta > 2$, using the Pareto prior from b) above. If we lose \$1 for choosing H_0 when in fact H_1 is true, and we lose \$ w for choosing H_1 when in fact H_0 is true, what decision $\delta(\mathbf{x})$ ($= 0$ or 1) should we make to minimize our expected loss? This may depend on w , of course... Show your work.

$$\delta(\mathbf{x}) =$$

Problem 5: Part (a) of this problem is completely unrelated to parts (b) and (c).

- a) (10) In a famous experiment, Gregor Mendel observed the shape and color of peas that resulted from certain crossbreedings. The reported results of one such experiment were:

round yellow	315	wrinkled yellow	101
round green	108	wrinkled green	32

According to his theory, the frequencies of these four classifications should be in the ratio 9 : 3 : 3 : 1. What do you conclude from a χ^2 test? Please give the value “ χ^2 ” of the test statistic (called “ Q ” by DeGroot) used for χ^2 tests, its number of degrees of freedom “d.f.”, and the P -value for the hypothesis that Mendel’s theory is correct, “ P .”

$\chi^2 =$ _____ d.f. = _____ $P =$ _____

Problem 5 (cont):

- b) (5) The 10 random variables $\{X_i\}$ are independent and take nonnegative integer values. Describe how you would use a generalized likelihood ratio test (GLRT) to test the hypothesis H_0 that these come from a Poisson distribution with mean $\lambda = 3$ against the composite alternative H_1 that they come from a Poisson distribution with some other (unspecified) mean. What statistics would you need from the data? Give a test statistic $T(\mathbf{x})$ such that your recommended test would reject H_0 for large values of T .

$$H_0 : \{X_i\}_{i \leq n} \sim \text{Po}(\lambda) \quad \text{for } \lambda = 3 \quad \text{vs.}$$

$$H_1 : \{X_i\}_{i \leq n} \sim \text{Po}(\lambda) \quad \text{for } \lambda \neq 3.$$

- (xc) Find the approximate probability distribution of T , if H_0 is true:

Problem 5 (cont):

- c) (5) The 10 random variables $\{X_i\}$ are (still) independent and take nonnegative integer values. Describe how you would use a GLRT to test the hypothesis H_0 that $\{X_i\}$ come from a Poisson distribution (with unspecified mean λ) against the alternative H_1 that they come from a Binomial distribution with $n = 5$ (but p unspecified):

$$H_0 : \{X_i\}_{i \leq n} \sim \text{Po}(\lambda) \quad \text{for some } \lambda \in [0, \infty) \quad \text{vs.}$$


$$H_1 : \{X_i\}_{i \leq n} \sim \text{Bi}(5, p) \quad \text{for some } p \in [0, 1]$$

What statistics would you need from the data? Give a test statistic $T(\mathbf{x})$ such that your recommended test would reject H_0 for large values of T .

Name: _____

Mth 136 = Sta 114

Extra worksheet, if needed:




Area $\Phi(x)$ under the Standard Normal Curve to the left of x .

x	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998

$\Phi(0.6745) = 0.75$ $\Phi(1.6449) = 0.95$ $\Phi(2.3263) = 0.99$ $\Phi(3.0902) = 0.999$
 $\Phi(1.2816) = 0.90$ $\Phi(1.9600) = 0.975$ $\Phi(2.5758) = 0.995$ $\Phi(3.2905) = 0.9995$

Critical Values for χ^2


$$\alpha = \int_{\chi^2_{\alpha}}^{\infty} c x^{\nu/2-1} e^{-x/2} dx$$



ν	$\chi^2_{.50}$	$\chi^2_{.25}$	$\chi^2_{.10}$	$\chi^2_{.05}$	$\chi^2_{.025}$	$\chi^2_{.01}$	$\chi^2_{.005}$	$\chi^2_{.001}$	$\chi^2_{.0005}$	$\chi^2_{.0001}$
1	0.4549	1.3233	2.7055	3.8415	5.0239	6.6349	7.8794	10.8276	12.1157	15.1367
2	1.3863	2.7726	4.6052	5.9915	7.3778	9.2103	10.5966	13.8155	15.2018	18.4207
3	2.3660	4.1083	6.2514	7.8147	9.3484	11.3449	12.8382	16.2662	17.7300	21.1075
4	3.3567	5.3853	7.7794	9.4877	11.1433	13.2767	14.8603	18.4668	19.9974	23.5127
5	4.3515	6.6257	9.2364	11.0705	12.8325	15.0863	16.7496	20.5150	22.1053	25.7448
6	5.3481	7.8408	10.6446	12.5916	14.4494	16.8119	18.5476	22.4577	24.1028	27.8563
7	6.3458	9.0371	12.0170	14.0671	16.0128	18.4753	20.2777	24.3219	26.0178	29.8775
8	7.3441	10.219	13.3616	15.5073	17.5345	20.0902	21.9550	26.1245	27.8680	31.8276
9	8.3428	11.389	14.6837	16.9190	19.0228	21.6660	23.5894	27.8772	29.6658	33.7199
10	9.3418	12.549	15.9872	18.3070	20.4831	23.2092	25.1882	29.5883	31.4198	35.5640
11	10.341	13.701	17.2750	19.6751	21.9200	24.7249	26.7568	31.2641	33.1366	37.3670
12	11.340	14.845	18.5493	21.0260	23.3366	26.2169	28.2995	32.9095	34.8213	39.1344
13	12.340	15.984	19.8119	22.3620	24.7356	27.6882	29.8195	34.5282	36.4778	40.8707
14	13.339	17.117	21.0641	23.6848	26.1189	29.1412	31.3193	36.1233	38.1094	42.5793
15	14.339	18.245	22.3071	24.9958	27.4884	30.5779	32.8013	37.6973	39.7188	44.2632
16	15.338	19.369	23.5418	26.2962	28.8453	31.9999	34.2672	39.2524	41.3081	45.9249
17	16.338	20.489	24.7690	27.5871	30.1910	33.4087	35.7185	40.7902	42.8792	47.5664
18	17.338	21.605	25.9894	28.8693	31.5264	34.8053	37.1565	42.3124	44.4338	49.1894
19	18.338	22.718	27.2036	30.1435	32.8523	36.1909	38.5823	43.8202	45.9731	50.7955
20	19.337	23.828	28.4120	31.4104	34.1696	37.5662	39.9968	45.3147	47.4985	52.3860
21	20.337	24.935	29.6151	32.6706	35.4789	38.9322	41.4011	46.7970	49.0108	53.9620
22	21.337	26.039	30.8133	33.9244	36.7807	40.2894	42.7957	48.2679	50.5111	55.5246
23	22.337	27.141	32.0069	35.1725	38.0756	41.6384	44.1813	49.7282	52.0002	57.0746
24	23.337	28.241	33.1962	36.4150	39.3641	42.9798	45.5585	51.1786	53.4788	58.6130
25	24.337	29.339	34.3816	37.6525	40.6465	44.3141	46.9279	52.6197	54.9475	60.1403
26	25.336	30.435	35.5632	38.8851	41.9232	45.6417	48.2899	54.0520	56.4069	61.6573
27	26.336	31.528	36.7412	40.1133	43.1945	46.9629	49.6449	55.4760	57.8576	63.1645
28	27.336	32.620	37.9159	41.3371	44.4608	48.2782	50.9934	56.8923	59.3000	64.6624
29	28.336	33.711	39.0875	42.5570	45.7223	49.5879	52.3356	58.3012	60.7346	66.1517
30	29.336	34.800	40.2560	43.7730	46.9792	50.8922	53.6720	59.7031	62.1619	67.6326
40	39.336	45.616	51.8051	55.7585	59.3417	63.6907	66.7660	73.4020	76.0946	82.0623
50	49.335	56.334	63.1671	67.5048	71.4202	76.1539	79.4900	86.6608	89.5605	95.9687
60	59.335	66.981	74.3970	79.0819	83.2977	88.3794	91.9517	99.6072	102.695	109.503
70	69.335	77.577	85.5270	90.5312	95.0232	100.425	104.215	112.317	115.578	122.755
80	79.334	88.130	96.5782	101.879	106.629	112.329	116.321	124.839	128.261	135.782
90	89.334	98.650	107.565	113.145	118.136	124.116	128.299	137.208	140.782	148.627
100	99.334	109.14	118.498	124.342	129.561	135.807	140.169	149.449	153.167	161.319

Critical Values for Student's t

$$p = \int_{-\infty}^{t_p} c \frac{dt}{(1 + t^2/\nu)^{(\nu+1)/2}}$$



ν	$t_{.60}$	$t_{.70}$	$t_{.80}$	$t_{.85}$	$t_{.90}$	$t_{.95}$	$t_{.975}$	$t_{.99}$	$t_{.995}$	$t_{.999}$	$t_{.9995}$	$t_{.9999}$
1	0.325	0.727	1.376	1.9626	3.078	6.314	12.76	31.82	63.66	318.3	636.6	3183.
2	0.289	0.617	1.061	1.3862	1.886	2.920	4.303	6.965	9.925	22.33	31.60	70.70
3	0.277	0.584	0.978	1.2498	1.638	2.353	3.182	4.541	5.841	10.22	12.92	22.20
4	0.271	0.569	0.941	1.1896	1.533	2.132	2.776	3.747	4.604	7.173	8.610	13.03
5	0.267	0.559	0.920	1.1558	1.476	2.015	2.571	3.365	4.032	5.893	6.869	9.678
6	0.265	0.553	0.906	1.1342	1.440	1.943	2.447	3.143	3.707	5.208	5.959	8.025
7	0.263	0.549	0.896	1.1192	1.415	1.895	2.365	2.998	3.499	4.785	5.408	7.063
8	0.262	0.546	0.889	1.1081	1.397	1.860	2.306	2.896	3.355	4.501	5.041	6.442
9	0.261	0.543	0.883	1.0997	1.383	1.833	2.262	2.821	3.250	4.297	4.781	6.010
10	0.260	0.542	0.879	1.0931	1.372	1.812	2.228	2.764	3.169	4.144	4.587	5.694
11	0.260	0.540	0.876	1.0877	1.363	1.796	2.201	2.718	3.106	4.025	4.437	5.453
12	0.259	0.539	0.873	1.0832	1.356	1.782	2.179	2.681	3.055	3.930	4.318	5.263
13	0.259	0.538	0.870	1.0795	1.350	1.771	2.160	2.650	3.012	3.852	4.221	5.111
14	0.258	0.537	0.868	1.0763	1.345	1.761	2.145	2.624	2.977	3.787	4.140	4.985
15	0.258	0.536	0.866	1.0735	1.341	1.753	2.131	2.602	2.947	3.733	4.073	4.880
16	0.258	0.535	0.865	1.0711	1.337	1.746	2.120	2.583	2.921	3.686	4.015	4.791
17	0.257	0.534	0.863	1.0690	1.333	1.740	2.110	2.567	2.898	3.646	3.965	4.714
18	0.257	0.534	0.862	1.0672	1.330	1.734	2.101	2.552	2.878	3.610	3.922	4.648
19	0.257	0.533	0.861	1.0655	1.328	1.729	2.093	2.539	2.861	3.579	3.883	4.590
20	0.257	0.533	0.860	1.0640	1.325	1.725	2.086	2.528	2.845	3.552	3.85	4.539
21	0.257	0.532	0.859	1.0627	1.323	1.721	2.080	2.518	2.831	3.527	3.819	4.493
22	0.256	0.532	0.858	1.0614	1.321	1.717	2.074	2.508	2.819	3.505	3.792	4.452
23	0.256	0.532	0.858	1.0603	1.319	1.714	2.069	2.500	2.807	3.485	3.768	4.415
24	0.256	0.531	0.857	1.0593	1.318	1.711	2.064	2.492	2.797	3.467	3.745	4.382
25	0.256	0.531	0.856	1.0584	1.316	1.708	2.060	2.485	2.787	3.450	3.725	4.352
26	0.256	0.531	0.856	1.0575	1.315	1.706	2.056	2.479	2.779	3.435	3.707	4.324
27	0.256	0.531	0.855	1.0567	1.314	1.703	2.052	2.473	2.771	3.421	3.690	4.299
28	0.256	0.530	0.855	1.0560	1.313	1.701	2.048	2.467	2.763	3.408	3.674	4.275
29	0.256	0.530	0.854	1.0553	1.311	1.699	2.045	2.462	2.756	3.396	3.659	4.254
30	0.256	0.530	0.854	1.0547	1.310	1.697	2.042	2.457	2.750	3.385	3.646	4.234
40	0.255	0.529	0.851	1.0500	1.303	1.684	2.021	2.423	2.704	3.307	3.551	4.094
60	0.254	0.527	0.848	1.0455	1.296	1.671	2.000	2.390	2.660	3.232	3.460	3.962
120	0.254	0.526	0.845	1.0409	1.289	1.658	1.980	2.358	2.617	3.160	3.373	3.837
∞	0.253	0.524	0.842	1.0364	1.282	1.645	1.960	2.326	2.576	3.090	3.291	3.719

Name	Notation	pdf/pmf	Range	Mean μ	Variance σ^2
Beta	$\text{Be}(\alpha, \beta)$	$f(x) = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1} (1-x)^{\beta-1}$	$x \in (0, 1)$	$\frac{\alpha}{\alpha+\beta}$	$\frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$
Binomial	$\text{Bi}(n, p)$	$f(x) = \binom{n}{x} p^x q^{(n-x)}$	$x \in 0, \dots, n$	np	$npq \quad (q = 1 - p)$
Exponential	$\text{Ex}(\lambda)$	$f(x) = \lambda e^{-\lambda x}$	$x \in \mathbb{R}_+$	$1/\lambda$	$1/\lambda^2$
Gamma	$\text{Ga}(\alpha, \lambda)$	$f(x) = \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x}$	$x \in \mathbb{R}_+$	α/λ	α/λ^2
Geometric	$\text{Ge}(p)$	$f(x) = p q^x$ $f(y) = p q^{y-1}$	$x \in \mathbb{Z}_+$ $y \in \{1, \dots\}$	q/p $1/p$	$q/p^2 \quad (q = 1 - p)$ $q/p^2 \quad (y = x + 1)$
HyperGeo.	$\text{HG}(n, A, B)$	$f(x) = \frac{\binom{A}{x} \binom{B}{n-x}}{\binom{A+B}{n}}$	$x \in 0, \dots, n$	nP	$nP(1-P) \frac{N-n}{N-1} \quad (P = \frac{A}{A+B})$
Logistic	$\text{Lo}(\mu, \beta)$	$f(x) = \frac{e^{-(x-\mu)/\beta}}{\beta[1+e^{-(x-\mu)/\beta}]^2}$	$x \in \mathbb{R}$	μ	$\pi^2 \beta^2 / 3$
Log Normal	$\text{LN}(\mu, \sigma^2)$	$f(x) = \frac{1}{x\sqrt{2\pi\sigma^2}} e^{-(\log x - \mu)^2 / 2\sigma^2}$	$x \in \mathbb{R}_+$	$e^{\mu + \sigma^2 / 2}$	$e^{2\mu + \sigma^2} (e^{\sigma^2} - 1)$
Neg. Binom.	$\text{NB}(\alpha, p)$	$f(x) = \binom{x+\alpha-1}{x} p^\alpha q^x$ $f(y) = \binom{y-1}{y-\alpha} p^\alpha q^{y-\alpha}$	$x \in \mathbb{Z}_+$ $y \in \{\alpha, \dots\}$	$\alpha q / p$ α / p	$\alpha q / p^2 \quad (q = 1 - p)$ $\alpha q / p^2 \quad (y = x + \alpha)$
Normal	$\text{No}(\mu, \sigma^2)$	$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\mu)^2 / 2\sigma^2}$	$x \in \mathbb{R}$	μ	σ^2
Pareto	$\text{Pa}(\alpha, \epsilon)$	$f(x) = \alpha \epsilon^\alpha / x^{\alpha+1}$	$x \in (\epsilon, \infty)$	$\frac{\epsilon \alpha}{\alpha-1}$	$\frac{\epsilon^2 \alpha}{(\alpha-1)^2 (\alpha-2)}$
Poisson	$\text{Po}(\lambda)$	$f(x) = \frac{\lambda^x}{x!} e^{-\lambda}$	$x \in \mathbb{Z}_+$	λ	λ
Snedecor F	$F(\nu_1, \nu_2)$	$f(x) = \frac{\Gamma(\frac{\nu_1+\nu_2}{2}) (\nu_1/\nu_2)^{\nu_1/2}}{\Gamma(\frac{\nu_1}{2}) \Gamma(\frac{\nu_2}{2})} \times$ $x^{\frac{\nu_1-2}{2}} \left[1 + \frac{\nu_1}{\nu_2} x \right]^{-\frac{\nu_1+\nu_2}{2}}$	$x \in \mathbb{R}_+$	$\frac{\nu_2}{\nu_2-2}$	$\left(\frac{\nu_2}{\nu_2-2} \right)^2 \frac{2(\nu_1+\nu_2-2)}{\nu_1(\nu_2-4)}$
Student t	t_ν	$f(x) = \frac{\Gamma(\frac{\nu+1}{2})}{\Gamma(\frac{\nu}{2}) \sqrt{\pi\nu}} [1 + x^2/\nu]^{-(\nu+1)/2}$	$x \in \mathbb{R}$	0	$\nu/(\nu-2)$
Uniform	$\text{Un}(a, b)$	$f(x) = \frac{1}{b-a}$	$x \in (a, b)$	$\frac{a+b}{2}$	$\frac{(b-a)^2}{12}$
Weibull	$\text{We}(\alpha, \beta)$	$f(x) = \alpha \beta x^{\alpha-1} e^{-\beta x^\alpha}$	$x \in \mathbb{R}_+$	$\frac{\Gamma(1+\alpha^{-1})}{\beta^{1/\alpha}}$	$\frac{\Gamma(1+2/\alpha) - \Gamma^2(1+1/\alpha)}{\beta^{2/\alpha}}$