

# STA732

## Statistical Inference

Lecture 09: Bayesian estimation

---

Yuansi Chen

Spring 2022

Duke University

<https://www2.stat.duke.edu/courses/Spring22/sta732.01/>



1. Construct minimum risk equivariant (MRE) estimator via conditioning on maximal invariant statistics
2. Pitman estimator of location
3. MRE for location is unbiased under squared error loss
4. MRE usually admissible

- We have finished the first approach of arguing for “the best” estimator in point estimation: by restricting to a small set of estimators
  - Unbiased estimators
  - Equivariant estimators
- We begin the second approach: global measure of optimality
  - average risk
  - minimax risk

1. Bayes risk, Bayes estimator
2. Examples
3. Bayes estimators are usually biased
4. Bayes estimators are usually admissible

Chap. 7 in Keener or Chap. 4 in Lehmann and Casella

## Bayes risk, Bayes estimator

---

## Recall the components of a decision problem

- Data  $X$
- Model family  $\mathcal{P} = \{P_\theta : \theta \in \Omega\}$ , a collection of probability distributions on the sample space
- Loss function  $L$ ,  $L(\theta, d)$  measures the loss incurred by the decision  $d$  when compared with the parameter obtained from  $\theta$
- Risk function  $R$ ,  $R(\theta, \delta) = \mathbb{E}_\theta[L(\theta, \delta)]$

## Motivation

It is in general hard to find uniformly minimum risk estimator. Oftentimes, we have risks that cross. This difficulty will not arise if the performance is measured via a single number.

## Def. Bayes risk

The **Bayes risk** is the average-case risk, integrated w.r.t. some measure  $\Lambda$ , called **prior**.

# The frequentist motivation of the Bayesian setup

## Motivation

It is in general hard to find uniformly minimum risk estimator. Oftentimes, we have risks that cross. This difficulty will not arise if the performance is measured via a single number.

## Def. Bayes risk

The **Bayes risk** is the average-case risk, integrated w.r.t. some measure  $\Lambda$ , called **prior**.

## Remark

For now, assume  $\Lambda(\Omega) = 1$  ( $\Lambda$  is a prob measure). Later we might deal with improper prior.



$$\begin{aligned}R_{\text{Bayes}}(\Lambda, \delta) &= \int_{\Omega} R(\theta, \delta) d\Lambda(\theta) \\ &= \mathbb{E}R(\Theta, \delta)\end{aligned}$$

where  $\Theta$  is the random variable with distribution  $\Lambda$ .

$$\mathbb{E}R(\Theta, \delta) = \mathbb{E}[\mathbb{E}[L(\Theta, \delta(X)) \mid X]]$$

Both  $X$  and  $\Theta$  are considered random.

The frequentist understanding: average risk makes sense without believing the parameter is random

An estimator  $\delta$  which minimizes the average risk  $R_{\text{Bayes}}(\Lambda, \cdot)$  is a **Bayes estimator**.

## Construct Bayes estimator

### Thm 7.1 in Keener

Suppose  $\Theta \sim \Lambda$ ,  $X \mid \Theta = \theta \sim P_\theta$ , and  $L(\theta, d) \geq 0$  for all  $\theta \in \Omega$  and all  $d$ . If

- $\mathbb{E}[L(\Theta, \delta_0)] < \infty$  for some  $\delta_0$
- for a.e.  $x$ , there exists a  $\delta_\Lambda(x)$  minimizing

$$\mathbb{E}[L(\Theta, d) \mid X = x]$$

with respect to  $d$

Then  $\delta_\Lambda$  is a Bayes estimator.

In words: the Bayes estimator can be found by minimizing the conditional distribution  $\mathbb{E}[L(\theta, d) \mid X = x]$ , one  $x$  at a time

## proof of Thm 7.1

## Def. Posterior

The conditional distribution of  $\Theta$  given  $X$ , written as  $\mathcal{L}(\Theta | X)$  is called the **posterior distribution**

## Remark

- $\Lambda$  is usually interpreted as prior belief about  $\Theta$  before seeing the data
- $\mathcal{L}(\Theta | X)$  is the belief after seeing the data

## Posterior calculation with density

Suppose prior density  $\lambda(\theta)$ , likelihood  $p_\theta(x)$ , then the posterior density is

$$\lambda(\theta | x) = \frac{\lambda(\theta)p_\theta(x)}{q(x)}$$

where  $q(x) = \int_{\Omega} \lambda(\theta)p_\theta(x)d\theta$  is the marginal density of  $X$ .

Then the Bayes estimator has the form

$$\delta_{\Lambda}(x) = \arg \min_d \int_{\Omega} L(\theta, d)\lambda(\theta | x)d\theta$$

## Posterior mean is Bayes estimator for squared error loss

Suppose  $L(\theta, d) = (g(\theta) - d)^2$  then the Bayes estimator is the posterior mean

proof:

## Examples

---



## Binomial model with Beta prior

Suppose  $X \mid \Theta = \theta \sim \text{Binomial}(n, \theta)$  with density  $\theta^x (1 - \theta)^{n-x} \binom{n}{x}$ ,  
 $\Theta \sim \text{Beta}(\alpha, \beta)$  with density  $\theta^{\alpha-1} (1 - \theta)^{\beta-1} \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$ . Find the  
Bayes estimator under squared error loss.

Suppose  $L(\theta, d) = w(\theta) (g(\theta) - d)^2$ . Find a Bayes estimator.

## Normal mean estimation

$$X \mid \Theta = \theta \sim \mathcal{N}(\theta, \sigma^2),$$

$$\Theta \sim \mathcal{N}(\mu, \tau^2).$$

Find the Bayes estimator of mean under squared error loss

What if we have  $n$  i.i.d. data points?

## Binary classification

Suppose the parameter space  $\Omega = \{0, 1\}$ .

$\mathbb{P}(X = x \mid \Theta = 0) = f_0(x)$  and  $\mathbb{P}(X = x \mid \Theta = 1) = f_1(x)$ . The prior is  $\pi(1) = p, \pi(0) = 1 - p$ .

Determine a Bayes estimator under 0-1 loss  $L(\theta, d) = \begin{cases} 0 & d = \theta \\ 1 & d \neq \theta \end{cases}$

**Bayes estimators are usually biased**

---

## Thm Lehmann Casella 4.2.3

If  $\delta$  is unbiased for  $g(\theta)$  with  $R_{\text{Bayes}}(\Lambda, \delta) < \infty$  then  $\delta$  is not Bayes under squared error loss unless its average risk is zero

$$\mathbb{E} [(\delta(X) - g(\Theta))^2] = 0$$

proof:

**Bayes estimators are usually  
admissible**

---



## Thm. Lehmann Casella 4.1.4

Let  $Q$  be the marginal distribution of  $X$ , i.e.,

$Q(E) = \int \mathbb{P}_\theta(E) d\Lambda(\theta)$ . Suppose  $L$  is strictly convex. If

1.  $R_{\text{Bayes}}(\Lambda, \delta_\Lambda) < \infty$ ,
2.  $Q(E) = 0$  implies  $P_\theta(E) = 0, \forall \theta$ ,

then the Bayes estimator  $\delta_\Lambda$  is unique.

proof: Use the following lemma

**Lem. Lehmann Casella exercise 1.7.26**

Let  $\phi$  be a strictly convex function over an interval  $I$ . If there exists a value  $a_0 \in I$  minimizing  $\phi$ , then  $a_0$  is unique.

### Thm. Lehmann Casella 5.2.4

A unique Bayes estimator (a.s. for all  $P_\theta$ ) is admissible.

proof:

- Bayes estimator is defined as the minimizer of the average risk over a prior on  $\theta$ .
- Bayes estimator can be constructed by conditioning the risk on each  $x$
- Bayes estimators are biased under squared error loss
- Bayes estimators are admissible under strictly convex loss

## What is next?

- Where do priors come from?
- Pros and cons of Bayes

Thank you for attending  
See you on Wednesday in Old  
Chem 025

