

# STA732

## Statistical Inference

Lecture 20: UMP restricted to unbiased tests

---

Yuansi Chen

Spring 2022

Duke University

<https://www2.stat.duke.edu/courses/Spring22/sta732.01/>



- Least favorable distributions as a way to reduce composite null to simple null

In general,  
think Lagrangian multiplier for constrained optimization  
think least favorable prior when dealing worst-case criteria!

1. What to do when UMP does not exist
2. General strategies for uniformly most powerful unbiased (UMPU) tests
3. UMP with power derivative restriction

Chap. 12.5-7 of Keener or Chap. 4 of Lehmann and Romano

## Beyond UMP

---

## Types of optimality:

Point estimation	Hypothesis testing
Uniform (in general does not exist) Restrict: UMVU, MRE Global: Bayes, Minimax Asymptotics	<b>UMP</b> Chap 4-6 in Lehmann and Romano Chap 8 Chap 11-13

## Restrict to smaller class of test functions

- Unbiased test
- Invariance
- Monotonicity

## Global measures

- Maximize the average power: put a prior on  $\Omega_1$
- Maximize worst case power: maximize the minimum power over  $\Omega_1$

### Def. Unbiased test, 12.25 in Keener

Let  $\alpha \in [0, 1]$ . A test  $\phi$  is unbiased level- $\alpha$  if

$$\beta_\phi(\theta) \leq \alpha, \forall \theta \in \Omega_0 \text{ and } \beta_\phi(\theta) \geq \alpha, \forall \theta \in \Omega_1$$

### Remark

- Unbiasedness enforces the appealing property that the probability of rejection is greater under any alternative distribution than it is under any null distribution
- It is related to a special case of risk unbiasedness if we design the loss function  $L$  such that  $L(\theta_0, \text{reject}) = 1 - \alpha$  and  $L(\theta_0, \text{accept}) = \alpha$ .

## Restrict test with some invariance (not covered)

Let  $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} \mathcal{N}(\theta, \sigma^2)$  for  $\sigma, \theta$  both unknown, and test  $H_0 : \theta = 0$  versus  $H_1 : \theta \neq 0$ . For  $i \in \{1, \dots, n\}$ , let  $X'_i = cX_i$  with  $c > 0$ . Then  $\mathbb{E}(X'_i) = \theta' = c\theta$ . Since testing  $\theta = 0$  is equivalent to testing  $\theta' = 0$ , it is natural to impose the invariance constraint

$$\forall c > 0 \quad \phi(X) = \phi(cX)$$

Such a test is unaffected by arbitrary rescaling of the data (which might occur when changing units from centimeters to meters).



## Restrict to tests with monotonicity (not covered)

Let  $X, Y$  be independent,  $X \sim \mathcal{N}(\theta_X, 1)$  and  $Y \sim \mathcal{N}(\theta_Y, 1)$  for  $\theta_X, \theta_Y$  unknown, and test  $H_0 : \theta_X \leq 0, \theta_Y \leq 0$ .

**A monotonicity restriction** requires that if  $\phi$  rejects upon observing  $(x, y)$ , then it should also reject for  $(x', y')$  where  $x' > x$  and  $y' > y$ .

## General strategies for UMPU

---

## Strategy outline

1. Prove that unbiasedness implies weaker constraints ( $\alpha$ -similarity)
2. Fix an alternative hypothesis
3. Find a MP test  $\phi$  under the weaker constraints (generalization of Neyman-Pearson lemma)
4. If  $\phi$  does not depend on the alternative hypothesis, then it is UMP for the composite alternative under the weaker constraints
5. Show  $\phi$  is UMP under the original constraint (unbiasedness).

Testing  $H_0 : \theta \in \Omega_0$  vs  $H_1 : \theta \in \Omega_1$ .  $\Omega_0$  and  $\Omega_1$  are subsets of a Euclidean space. Let  $\omega$  be the **common boundary** between  $\Omega_0$  and  $\Omega_1$ :

$$w = \bar{\Omega}_0 \cap \bar{\Omega}_1$$

In words,  $\omega$  is the intersection of the closures of  $\Omega_0$  and  $\Omega_1$

### Example 1

Testing  $H_0 : \theta = \theta_0$  vs  $H_1 : \theta \neq \theta_0$ , then  $\omega = \{\theta_0\}$

### Example 2

Testing  $H_0 : \theta_1 \leq \tilde{\theta}$  vs  $H_1 : \theta_1 > \tilde{\theta}$  in the presence of nuisance parameters  $(\theta_2, \dots, \theta_{k+1})$ , then

$$\omega = \{\theta \in \mathbb{R}^{k+1} : \theta_1 = \tilde{\theta}\}$$

**Def.  $\alpha$ -similarity, 4.1 in Lehmann and Romano**

A test  $\phi$  satisfying  $\mathbb{E}_\theta \phi(X) = \alpha$  for all  $\theta \in \omega$  is called  **$\alpha$ -similar** on  $\omega$

## **Relation to unbiasedness**

When  $\beta_\phi(\theta)$  is continuous in  $\theta$ , unbiasedness implies  $\alpha$ -similarity on  $\omega$ .

draw a picture

## Lem. 4.1.1 Lehmann and Romano

If  $\theta \mapsto \beta_\phi(\theta)$  is continuous on  $\Omega$  for all  $\phi$ , and  $\phi_0$  is a UMP test among  $\alpha$ -similar level- $\alpha$  tests, then  $\phi_0$  is also UMPU at level  $\alpha$

### Proof: compare to the constant test

$\phi_0$  is UMP among  $\alpha$ -similar tests, it is at least as powerful as the constant test  $\phi_\alpha(X) \equiv \alpha$ .

## UMPU in two-sided testing without nuisance params

Testing  $H_0 : \theta = \theta_0$  vs  $H_1 : \theta \neq \theta_0$ . Suppose  $X$  is from a 1-param exp family

$$p_{\theta}(x) = h(x) \exp(\theta T(x) - A(\theta))$$



## UMPU in two-sided testing without nuisance params

Testing  $H_0 : \theta = \theta_0$  vs  $H_1 : \theta \neq \theta_0$ . Suppose  $X$  is from a 1-param exp family

$$p_\theta(x) = h(x) \exp(\theta T(x) - A(\theta))$$

- We know that no UMP test exist in the Gaussian case
- Assume  $\phi$  is unbiased at level- $\alpha$ , then

$$\beta_\phi(\theta_0) = \mathbb{E}_{\theta_0} \phi(X) = \alpha$$

$$\beta_\phi(\theta_0) \leq \beta_\phi(\theta) \text{ for all } \theta \in \mathbb{R}$$

- If we further assume  $\beta_\phi$  is differentiable, then the second point translate to

$$0 = \beta'_\phi(\theta_0) = \int \phi(x) \frac{d}{d\theta} p_{\theta_0}(x) d\mu(x)$$

To find UMPU in two-sided testing, we first find UMP with power derivative constraint

$$\begin{aligned} \max_{\phi} \beta_{\phi}(\theta') \quad & \forall \theta' \in \Omega_1 \\ \text{s.t. } \beta_{\phi}(\theta_0) &= \alpha \\ \beta'_{\phi}(\theta_0) &= 0 \end{aligned}$$

Method of undetermined multipliers allow us to deal with UMP problems with multiple constraints!

## Proof for UMP with power derivative constraint in two-sided testing, 1-param exp family

- Fix a simple alternative  $\theta' > \theta_0$
- Use method of undetermined multipliers to determine a rejection region for the simple vs simple testing
- Discuss the shape of the rejection region
- Find UMP test for  $H_0 : \theta_0, H_1 : \theta' > \theta_0$
- Reverse the above argument to show the same test works for  $H_0 : \theta_0, H_1 : \theta' < \theta_0$
- The test does not depend on the alternative, so UMP for the composite alternative

## Recall: Methods of Undetermined Multipliers applied to testing (1)

We plan to apply the Methods of Undetermined Multipliers to the case  $U$  is the space of test functions  $\phi$ :

$$F_i(\phi) = \int \phi(x) f_i(x) d\mu(x).$$

We want to

$$\begin{aligned} \max \quad & \int \phi(x) f_{m+1}(x) d\mu(x) \\ \text{s.t.} \quad & \int \phi(x) f_i(x) d\mu(x) = c_i, \quad \forall i = 1, \dots, m \end{aligned}$$

## Recall: Methods of Undetermined Multipliers applied to testing (2)

According to Lem 3.6.1, we consider to maximize

$$F_{m+1}(\phi) - \sum_i k_i F_i(\phi) = \int \phi(x) \left( f_{m+1}(x) - \sum_{i=1}^m k_i f_i(x) \right) d\mu(x)$$

It is not hard to show (ignoring all regularity assumptions), the optimal solution should have the form

$$\phi(x) = \begin{cases} 1 & \text{if } f_{m+1}(x) > \sum_{i=1}^m k_i f_i(x) \\ 0 & \text{if } f_{m+1}(x) < \sum_{i=1}^m k_i f_i(x) \end{cases}$$

Finally, we choose  $k_i$  so that the constraints are all satisfied

Existence of  $\phi^*$  in general space (convex and closed) requires some technical details, see Chapter 12.5 Keener

## Conclude that the UMP test with with power derivative constraint is also UMPU

- First,  $\phi_\alpha \equiv \alpha$  also satisfies the two constraints. Since  $\phi$  is more powerful, then  $\phi$  is unbiased.
- Second, all unbiased tests satisfy the two constraints.  
Conclude that  $\phi$  is UMP among all level- $\alpha$  unbiased tests.

## Example

Suppose  $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, \sigma^2)$  with  $H_0 : \sigma = \sigma_0$  vs  $H_1 : \sigma \neq \sigma_0$

- UMPU exists in two sided testing without nuisance parameters
- UMPU via method of undetermined multipliers



### UMPU in multiparameter exp family

- Nuisance parameters
- The idea of conditioning

Thank you for attending  
See you on Monday in Old Chem  
025

